

# Asian Journal of Natural Product Biochemistry

| Asian J Nat Prod Biochem | vol. 20 | no. 1 | June 2022 |  
| ISSN 2775-4189 | E-ISSN 2775-4197 |

Palm sugar (*Arenga pinata*) photo by Grid.id

# Asian Journal of Natural Product Biochemistry

| Asian J Nat Prod Biochem | vol. 20 | no. 1 | June 2022 |

---

<b>Biosafety evaluation of ethanolic extract of <i>Phyllanthus amarus</i> leaves on liver and kidney of wistar rats</b> OJOCHENEMI EJEH YAKUBU, MICHAEL SUNDAY ABU, OKONKWO CHIBUIKE INNOCENT, JOB ITANYI ONUCHE	1-5
<b>Comparison of phytochemical constituents of ethanol leaf extracts of <i>Solanum macrocarpon</i> and <i>Vernonia amygdalina</i></b> OLUSOJI ADEBUSOYE OYESOLA, IQUOT ISAAC SAMPSON, ADEBIYI ADELOWO AUGUSTINE, OLUKADE BALIQIS ADEJOKE, GEORGE EMMANUEL TAIWO	6-10
<b>The effect of palm sap (<i>Arenga pinnata</i>) on blood glucose levels of mice (<i>Mus musculus</i>) alloxan-induced diabetic</b> SINDI MARYAM LIPUTO, DJUNA LAMONDO, MARGARETHA SOLANG	11-15
<b>Short Communication: Extraction and characterization of pectin from ripe and unripe mango (<i>Mangifera indica</i>) peel</b> CHRISTOPHER O. SHAIBU, JULIUS DINSHIYA, VIVIAN EBERE SHAIBU	16-20
<b>Extraction, purification, and quantification of hesperidin from the immature <i>Citrus grandis</i> maxima fruit Nepal cultivar</b> HEMANTH KUMAR MANIKYAM, PRATHIBHA TRIPATHI, SANDEEP BALVANT PATIL, JANARDAN LAMICHHANE, MVNL CHAITANYA, ABHINANDAN RAVSAHEB PATIL	21-26



# Asian Journal of Natural Product Biochemistry

| Asian J Nat Prod Biochem | vol. 20 | no. 1 | June 2022 |

## ONLINE

<http://smujo.id/jnpb>

## p-ISSN

2775-4189

## e-ISSN

2775-4197

## PUBLISHER

Society for Indonesian Biodiversity

## CO-PUBLISHER

Universitas Sebelas Maret, Surakarta, Indonesia

## OFFICE ADDRESS

Jl. Ir. Sutami 36A, Surakarta 57126, Central Java, Indonesia. Tel./fax. +62-271-663375, email: [editors@smujo.id](mailto:editors@smujo.id)

## PERIOD OF ISSUANCE

June, December

## EDITOR-IN-CHIEF

**Khalid Ahmad Khalid** – National Research Center, Cairo, Egypt

## ASSOCIATE EDITOR

**Ahmad Dwi Setyawan** – Universitas Sebelas Maret, Surakarta, Indonesia

## EDITORIAL BOARD

**Andri Frediansyah** – Unit for Natural Product Technology, National Research and Innovation Agency, Indonesia

**Asish Kumar Parida** – CSIR-Central Salt & Marine Chemicals Research Institute, Bhavnagar, India

**C.J. Sugiharjo** – Universitas Gadjah Mada, Yogyakarta, Indonesia

**Cedric B. Baker** – Mercer University, Atlanta, USA

**Chaerul** – Research Center for Biology, National Research and Innovation Agency, Indonesia

**Dayar Arbain** – Universitas Andalas Padang, Indonesia

**Enos Tangke Arung** – Universitas Mulawarman, Samarinda, Indonesia

**M. Ja'far Luthfi** – UIN Sunan Kalijaga, Yogyakarta, Indonesia

**Mallappa Kumara Swamy** – Universiti Putra Malaysia, Malaysia

**Supriyadi** – Indonesian Spices and Medicinal Crops Research Institute, Bogor, Indonesia

**Suranto** – Universitas Sebelas Maret, Surakarta, Indonesia

**Syamsul Arifin Achmad** – Institut Teknologi Bandung, Indonesia

**Taufik Hidayat** – Institut Pertanian Bogor, Indonesia

*List of reviewers: <https://smujo.id/jnpb/reviewers>*



Society for Indonesia  
Biodiversity



Universitas Sebelas Maret  
Surakarta, Indonesia

## GUIDANCE FOR AUTHORS

**Aims and Scope** *Asian Journal of Natural Product Biochemistry* (**Asian J Nat Prod Biochem**) (formerly *Biofarmasi*) encourages submission of manuscripts dealing with all important contributions to natural product chemistry, include bioactivity compounds of land and sea and of plants, microbes and animals; activities of antimicrobial, antiviral, antiparasite, anti-cancer, etc.; ethnopharmacology, ethnomedicines and medical application; plant-derived drugs and pharmaceutical formulations and evaluation; as well as biochemistry, chemistry, biotechnology, physiology, cell structures, pharmacology, toxicology, and chemical ecology of natural products.

**Article types** The journal seeks for: (i) **Research papers**, (ii) **Reviews**, and (iii) **Short communications**. Original full-length research manuscripts are limited to 8,000 words (including tables and figures) or proportional to articles in this publication number (beyond that, it should be with notice). Review articles are also limited to 8,000 words, while Short communications should be less than 2,500 words, except for pre-study (can be more).

**Submission:** The journal only accepts online submissions through the open journal system (<https://smujo.id/jnpb/about/submissions>) or, for login problems, email the editors at [unsjournals@gmail.com](mailto:unsjournals@gmail.com) (or [editors@smujo.id](mailto:editors@smujo.id)). Submitted manuscripts should be the original works of the author(s). Please ensure that the manuscript is submitted using the template, which can be found at (<https://biodiversitas.mipa.uns.ac.id/D/template.doc>). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, and a paragraph describing the claimed novelty of the findings versus current knowledge. Please also provide a list of five potential reviewers in your cover letter. They should come from outside your institution and better from three different countries. Submission of a manuscript implies the submitted work has not been published (except as part of a thesis or report, or abstract) and is not being considered for publication elsewhere. When a group writes a manuscript, all authors should read and approve the final version of the submitted manuscript and its revision; and agree on the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis, drafting the manuscript, and correcting the revision. All authors must be responsible for the work's quality, accuracy, and ethics.

**Ethics** Author(s) must be obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

**Copyright** If the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. For the new invention, authors must manage its patent before publication.

**Open Access** The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

**Acceptance** Only articles written in US English are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double-blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. Manuscripts will be rejected if the content does not align with the journal scope, does not meet the standard quality, is in an inappropriate format, or contains complicated grammar, dishonesty (i.e., plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and significance. **Uncorrected proofs** will be sent to the corresponding author by system or email as .doc or .docx files for checking and correcting typographical errors. The corrected proofs should be returned in 7 days to avoid publication delays. The accepted papers will be published online in chronological order at any time but printed at the end of each month.

**Free of charge** This publication is dedicated entirely to the advancement of science and technology, therefore author(s), or author institution(s) are not subject to publication fees. **Reprints** The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by email and sending back the uncorrected proofs.

**Manuscript preparation** Manuscript is typed on A4 (210x297 mm<sup>2</sup>) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering sizes can be applied in presenting tables and figures (9 pt). Word processing program or additional software can be used; however, it must be PC compatible, use the template, and use Microsoft Word based (.doc or .rtf; not .docx). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italics, except in italicized sentences. Scientific names (genus, species, author) and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. The genus name can be shortened after the first mention, except in early sentences, or where this may generate confusion; name of the author can be eliminated after the first mention. For example, *Rhizopus oryzae* L. UICC 524 can be written hereinafter as *R. oryzae* UICC 524. Using trivial names should be avoided. **Biochemical and chemical nomenclature** should follow the order of the IUPAC-IUB. For DNA sequences, it is better to use Courier New font. Standard chemical abbreviations can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurements** should use IS denominations, and other systems should use equivalent values with the denomination of IS mentioned first. A dot should not follow abbreviations like g, mg, mL, etc. Minus index (m<sup>-2</sup>, L<sup>-1</sup>, h<sup>-1</sup>) suggested being used, except in things like "per-plant" or "per-plot." **Mathematical equations** can be written down in one column with text; in that case, they can be written separately. **Numbers** one to ten are written in words, except if it relates to measurement, while values above them are written in number, except in early sentences. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent." Avoid expressing ideas with complicated sentences and verbiage/phrasing, and use efficient and effective sentences.

**The title** of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Name of author(s) should be completely written, especially for the first and the last name. **Name and institution** address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. We choose local names in Bahasa Indonesia for universities in Indonesia. The mention of "strata" program, should be avoided. Manuscript written by a group, author for correspondence along with address is required (marked with "\*"). **The title page** (first page) should include title of the article, full name(s), institution(s) and address(es) of the author(s); the corresponding authors detailed postage and e-mail addresses (P), and phone (O) and fax numbers (O).

**Abstract** A concise abstract is required (about 200 words). The abstract should be informative and state briefly the aim of the research, the principal results and major conclusions. An abstract is often presented separately from the article, thus it must be able to stand alone (completely self-explanatory). References should not be cited, but if essential, then cite the author(s) and year(s). Abbreviations should be avoided, but if essential, they must be defined at their first mention. **Keywords** are about five words, covering scientific and local name (if any), research themes, and special methods used; and sorted from A to Z. **Abbreviations** (if any): All important abbreviations must be defined at their first mention there. **Running title** is about five words.

**Introduction** is about 600 words, covering the aims of the research and provide an adequate background, avoiding a detailed literature survey or a summary of the results. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for a manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. **Concluding** sentence should be given at the end of the discussion. **Acknowledgements** are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest are noted.

**Figures and Tables** of a maximum of three pages should be clearly presented. The title of a picture is written down below the picture, while the title of a table is written above the table. Colored figures can only be accepted if the information in the manuscript can lose without those images; the chart is preferred to use black and white images. The author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned the source. Author is suggested referring to Wikipedia for international boundaries and Google Earth for satellite imagery. If not specifically mentioned, it is assumed to refer to these sources. **There is no appendix**, all data or data analysis is incorporated into Results and Discussions. For broad data, it can be displayed on the website as a supplement.

**References** Preferably 80% of it comes from scientific journals published in the last 10 years. In the text, give the author names followed by the year of publication and arrange from oldest to newest and from A to Z; in citing an article written by two authors, both of them should be mentioned; however, for three and more authors only the first author is mentioned followed by et al. For example, Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation should be avoided, as shown with the word "cit." Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in alphabetical order. Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** ([www.issn.org/2-22661-LTWA-online.php](http://www.issn.org/2-22661-LTWA-online.php)). Please include DOI links for journal papers. The following examples are for guidance.

### Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158. DOI: 10.13057/biodiv/d070213.

The usage of "et al." in long author lists will also be accepted:

Smith J, Jones M Jr, Houghton L et al. 1999. Future of health insurance. *N Engl J Med* 965: 325-329. DOI: 10.1007/s002149800025.

### Book:

Rai MK, Carpinella C. 2006. *Naturally Occurring Bioactive Compounds*. Elsevier, Amsterdam.

### Chapter in the book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds.). *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

### Abstract:

Assaeed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

### Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.). *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

### Thesis, Dissertation:

Sugiyarto. 2004. *Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon*. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

### Information from the internet:

Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4: 187. DOI: 10.1038/msb.2008.24. [www.molecularsystembiology.com](http://www.molecularsystembiology.com).

**THIS PAGE INTENTIONALLY LEFT BLANK**

## Biosafety evaluation of ethanolic extract of *Phyllanthus amarus* leaves on liver and kidney of wistar rats

OJOCHENEMI EJEH YAKUBU, MICHAEL SUNDAY ABU\*, OKONKWO CHIBUIKE INNOCENT, JOB ITANYI ONUCHE

Department of Biochemistry, Faculty of Pure and Applied Sciences, Federal University Wukari. PMB 1020 Katsina-Ala Road, Wukari, Taraba State, Nigeria. Tel./fax.: +23-480-000-0000, \*email: abulex77@yahoo.com

Manuscript received: 30 December 2021. Revision accepted: 20 March 2022.

**Abstract.** Yakubu OE, Abu MS, Innocent OC, Onuche JI. 2022. Biosafety evaluation of ethanolic extract of *Phyllanthus amarus* leaves on liver and kidney of wistar rats. *Asian J Nat Prod Biochem* 20: 1-5. Medicinal plants such as *Phyllanthus amarus* Schumach. & Thonn. have played a significant role in maintaining human health since ancient times. Documented evidence shows that the plant has been used to treat various ailments. Still, its biosafety on vital organs such as the liver and kidney has not been specifically studied. The study was designed to ascertain the biosafety of ethanolic extract of *P. amarus* leaves on the liver and kidney of wistar rats. A total of twenty (20) wistar rats were used, divided into 4 groups of 5 animals each (Group A, B, C, D). Group B, C, D received 100 mg/kg, 200 mg/kg, 400 mg/kg of the extract, respectively. Group A (Control) received only distilled water. The treatment was carried out for two weeks. At the end of the experiments, the animals were sacrificed. Afterward, liver function and renal function evaluations were conducted. The indicators of liver function include Total protein (TP), aspartate aminotransferase (AST), alkaline phosphatase (ALP), alanine aminotransferase (ALT), albumin (ALB), direct bilirubin (DB), and total bilirubin (TB) were determined. The indicators of renal function include sodium ion (Na<sup>+</sup>), potassium ion (K<sup>+</sup>), chlorine ion (Cl<sup>-</sup>), urea, and creatinine. The results showed that the parameters of the liver function of treated animals were no significant ( $p > 0.05$ ) difference compared to the normal rats (control). Concentrations of sodium ions (Na<sup>+</sup>), potassium ions (K<sup>+</sup>), chlorine ions (Cl<sup>-</sup>), urea, and creatinine in the treated animals were statistically comparable to the normal rats at ( $p < 0.05$ ). This research showed that *P. amarus* did not induce hepatotoxicity or nephrotoxicity at the dose of 100-400 mg/kg BW. Furthermore, this finding has validated the use of *P. amarus* as a traditional herb for treating and managing several diseases, including kidney and liver diseases, without significant organ damage.

**Keywords:** Biosafety, ethanolic extract, kidney, liver, *Phyllanthus amarus*, wistar rats

### INTRODUCTION

Medicinal plants have continued to play a dominant role in maintaining human health since ancient times. Most modern medicinal drugs are of plant origin (Tuhinadri and Sami 2015) and are essential in drug development programs of the pharmaceutical industry. The knowledge generated over decades has revealed medicinal plants such as *Phyllanthus amarus* Schumach. & Thonn. have been effective in the traditional management of various diseases. Based on these conditions, about 80% of the world population depends solely on traditional (Emeka et al. 2013) or herbal medicine to treat various diseases, especially in Africa and other developing nations (Ilonga et al. 2018; Yakubu et al. 2020a). However, the historical role of medicinal herbs in treating and preventing disease does not assure their safety for uncontrolled usage by an uninformed public (Matthews et al. 1999). Most medicinal plants have relatively non-toxic potential or adverse effects when used by humans; however, some medicinal plants have a high level of toxicity. In addition, they have the possibility of damaging organs. For this reason, caution in using medicinal plants is presently on the increase due to easy availability, affordability, accessibility, and promising efficacy compared to the often-high cost and adverse effects of the standard synthetic drug (Yakubu et al. 2020c).

*Phyllanthus amarus* is a broad-spectrum medicinal plant that has received worldwide recognition (Srividiya and Perival 1995). They are widely distributed in most tropical and subtropical countries and have long been used in traditional medicine to treat chronic liver diseases (Liu et al., 2003). *P. amarus* is generally employed to reduce pain, expel intestinal gas, stimulate and promote digestion, as anti-helminths to expel intestinal worms, and act as a mild laxative. *P. amarus* also has antiseptic, diuretic, antiviral, antidiabetic, hypotensive, and antipyretic, and it is also used to treat jaundice, diarrhea, dysentery, wound ulcers, and urogenital diseases (Calixto et al. 1998). The plant extracts have been found to contain high levels of saponins, tannins, flavonoids, and alkaloids (Fernand 1998; Naaz et al. 2007; Krithika and Verma 2009). *P. amarus* has been classified as having a low potential for toxicity, with an LD<sub>50</sub> averaging 2000 mg/kg/day (Krithika and Verma 2009). However, the toxicity of this plant on vital organs such as the liver and kidney needs to be revalidated, considering its increased demand for several herbal medicine preparations. Therefore, this study was designed to ascertain the biosafety of ethanolic extract of *P. amarus* leaves on the liver and kidney of wistar rats.

## MATERIALS AND METHODS

### Collection of plant

The fresh leaves of *P. amarus* were collected in December 2019 within the Biological garden of Federal University Wukari Taraba State, Nigeria. The plant was identified and authenticated in the herbarium of the Biological Science Department, Federal University Wukari, Nigeria. The fresh leaves were air-dried for three (3) days and pounded into a fine powder using a mortar and pestle.

### Extraction

The dried material (500 g) was macerated in 2000 mL of ethanol (1:4 w/v) for 48 hours at room temperature, according to Yakubu et al. (2014). It was continually stirred every 5 hours. After finishing maceration, the filtrate was sieved with clean white mesh before filtering using Whitman No 1 filter paper. Next, the filtrate was poured into a beaker and air-dried. Finally, the resultant extract (18 g) was placed into the incubator for further studies.

### Animals management

Twenty healthy adult wistar rats of both sexes were obtained and maintained in the animal house of the Biochemistry Department, Faculty of Pure and Applied Science, Federal University Wukari, Taraba State, Nigeria. They were acclimatized for two weeks and fed with growers' mash and water ad libitum.

### Experimental design

The wistar rats with an average weight of 177 g were randomly assigned into four (4) groups: A, B, C, and D of five (5) animals each. Group B, C, and D served as treatment groups, while group A served as the control. The rats in the treatment groups (B, C, and D) received 100 mg/kg, 200 mg/kg, and 400 mg/kg body weight of ethanolic extracts of *P. amarus* orally through an or gastric tube daily, respectively. The control group received an equal volume of distilled water without the *P. amarus* extract. After the experimental period, animals were sacrificed under chloroform anesthesia, venous blood was collected by cardiac puncture, and the liver was harvested. Blood samples were collected, and serum was obtained by centrifuging at 3000 rpm for 5 min.

Group A: Normal control received distilled water

Group B: 100 mg/kg/p.o/day of extract

Group C: 200 mg/kg/p.o/day of extract

Group D: 400 mg/kg/p.o/day of extract

### Determination of biochemical parameters

#### Analysis of Aspartate Aminotransferase (AST) activity

AST activity was determined by the method described by Amador and Wacker (1962).

**Principle:** L-Aspartate and  $\alpha$ -Ketoglutarate react in the presence of AST in the sample to yield oxaloacetate and L-glutamate. The oxaloacetate is reduced by malate dehydrogenase to yield L-malate with NADH oxidation to NAD<sup>+</sup>. The reaction is monitored by measuring the decrease in NADH absorbance at 340 nm. The reduction rate in absorbance is proportional to AST activity in the

sample.

**Procedure:** To 1 mL of reagent was added to all required test tubes, 0.05 mL of the sample was added to the sample test tube, and none to the blank. Next, it was incubated at room temperature for 20 minutes and mixed immediately, and the first absorbance of the test was read exactly at 1 minute and then at 30, 60, 90, and 120 seconds at 340 nm. Finally, the mean change in absorbance per minute was determined, and the test results were calculated as follows :

$$\text{Serum AST activity (IU/L)} = \text{Change in A/min} \times 3376.$$

#### Assessment of Alanine Aminotransferase (ALT) activity

ALT activity was determined by the method described by Amador and Wacker (1962).

**Principle:** L-alanine and  $\alpha$ -ketoglutarate react in the presence of ALT in the sample to yield Pyruvate and L-glutamate. Pyruvate is reduced by lactate dehydrogenase to yield lactate with oxidation of NADH to NAD. The reaction is monitored by measuring the decrease in absorbance at 340 nm. The rate of reduction is proportional to ALT activity in the sample.


**Procedure:** To 1 mL of reagent was added to all required test tubes, 0.05 mL of the sample was added to the test sample tube, and none to the blank. Next, it was incubated at room temperature for 20 min and mixed immediately, and the first absorbance was read at precisely 1 minute and then at 30, 60, 90, and 120 seconds at 340 nm. Finally, the mean change in absorbance per minute was determined, and test results were calculated as follows:

$$\text{Serum ALT activity (IU/L)} = \Delta A/\text{min} \times 3376.$$

#### Assessment of Alkaline Phosphatase (ALP) activity

The serum activity of Alkaline Phosphatase (ALP) was determined by the method Haussament (1977) described.

#### Principle:



Alkaline phosphatase in a sample hydrolyses para nitrophenyl phosphate into para nitrophenol and phosphate in the presence of magnesium ions. The absorbance increase rate of the reaction mixture at 405 nm and 37°C due to the liberation of para nitrophenol is proportional to the alkaline phosphatase activity.

**Procedure:** Reagent (1 mL) containing diethanolamine buffer, magnesium chloride, and substrate (P-nitrophenyl phosphate) was added into a clean test tube and incubated at 37°C, followed by the addition of 0.02 mL of sample. The mixture was mixed thoroughly, and the immediate absorbance of the sample was read precisely at 30, 60, 90, and 120 seconds at 405 nm against the reference blank (distilled water). The mean change in absorbance per minute was determined, and the test results were calculated.

The ALP activity was calculated using the following formulae:

Serum alkaline phosphatase activity (IU/L) =  $\Delta A/\text{min} \times 2713$ .

#### Serum creatinine and urea

Serum creatinine and urea were determined using an auto-analyzer: Selectra ProM, while the determination of Serum Electrolytes was carried out using the Ion-Selective Electrode (ISE) method (Burnett et al. 2000)

#### Statistical analysis

The mean  $\pm$  SD of all values was calculated, and the change observed between the treatment groups and control was subjected to analysis of variance (ANOVA) using SPSS version 23. Differences between groups were considered significant at  $p < 0.05$ .

## RESULTS AND DISCUSSION

#### Effect of *Phyllanthus amarus* leaves extract on serum electrolytes of wistar rat

The effect of the *P. amarus* leaves extract on some serum electrolyte levels is depicted in Table 1. There was no significant ( $p < 0.05$ ) difference between the rats administered with extract and the normal/control rats. However, a significant ( $p < 0.05$ ) decrease in the  $\text{Na}^+$  level of animals that received 100 and 200 mg/kg of the extract compared to the control animals. Similarly, a significant

( $p < 0.05$ ) reduction was observed in the  $\text{K}^+$  level of the group that received 100 mg/kg compared to the other extract groups, including the control group.

#### Effect of *Phyllanthus amarus* leaves extract on urea and creatinine level in wistar rat

All the groups, including the extract administered and the control groups, showed no significant ( $p > 0.05$ ) differences in the urea and creatinine levels, as represented in Table 2.

#### Effect of *Phyllanthus amarus* leaves extraction liver enzyme of wistar rat

Table 3 showed that the liver enzymes' activity was significantly ( $p < 0.05$ ) high or low in the group that received 200 mg/kg compared to the control group. However, the other extract-administered groups showed no consistent difference in enzyme activity compared with the control.

#### Effect of *Phyllanthus amarus* leaves extract on some liver function biomarkers of wistar rat

Rats treated with the extract at different doses did not show significant ( $p > 0.05$ ) differences in the total protein levels, total bilirubin, direct bilirubin, and albumin, as represented in Table 4 when compared to the control rats.

**Table 1.** Effect of *Phyllanthus amarus* leaves extract on serum electrolytes of wistar rat

Treatment	$\text{Na}^+$ (mmol/L)	$\text{K}^+$ (mmol/L)	CL (mmol/L)
Control	238.6 $\pm$ 1.34 <sup>b</sup>	38.7 $\pm$ 5.55 <sup>b</sup>	145.0 $\pm$ 25.14 <sup>a</sup>
Extract 100 mg	171.0 $\pm$ 06.74 <sup>a</sup>	36.18 $\pm$ 7.29 <sup>b</sup>	136.4 $\pm$ 19.06 <sup>a</sup>
Extract 200 mg	185.6 $\pm$ 05.85 <sup>a</sup>	19.5 $\pm$ 5.99 <sup>a</sup>	145.6 $\pm$ 19.96 <sup>a</sup>
Extract 400 mg	223.6 $\pm$ 26.15 <sup>b</sup>	39.7 $\pm$ 6.08 <sup>b</sup>	149.4 $\pm$ 19.93 <sup>a</sup>

Note: Results represent mean  $\pm$  standard deviation of group results (n=5). Values having different superscripts along the column are considered statistically significant ( $p < 0.05$ )

**Table 2.** Effect of *Phyllanthus amarus* leaves extract on urea and creatinine level in wistar rat

Treatment	Urea (mmol/L)	Creatinine (mmol/L)
Control	10.44 $\pm$ 5.01 <sup>a</sup>	5.12 $\pm$ 3.21 <sup>a</sup>
Extract 100 mg	0.7.84 $\pm$ 1.65 <sup>a</sup>	4.96 $\pm$ 3.68 <sup>a</sup>
Extract 200 mg	13.52 $\pm$ 5.67 <sup>a</sup>	3.12 $\pm$ 3.64 <sup>a</sup>
Extract 400 mg	12.68 $\pm$ 9.29 <sup>a</sup>	1.92 $\pm$ 2.10 <sup>a</sup>

Note: Results represent mean  $\pm$  standard deviation of group results (n=5). Values having different superscripts along the column are considered statistically significant ( $p < 0.05$ )

**Table 3.** Effect of *Phyllanthus amarus* leaves extraction liver enzyme of wistar rat

Treatment	ALT (IU/mL)	AST (IU/mL)	ALP (IU/L)
Control	3.8 $\pm$ 1.30 <sup>a</sup>	5.4 $\pm$ 0.54 <sup>a</sup>	16.0 $\pm$ 4.06 <sup>c</sup>
Extract 100 mg	11.4 $\pm$ 1.34 <sup>b</sup>	5.54 $\pm$ 3.13 <sup>a</sup>	13.2 $\pm$ 5.11 <sup>bc</sup>
Extract 200 mg	20.4 $\pm$ 6.46 <sup>c</sup>	10.4 $\pm$ 2.07 <sup>b</sup>	7.8 $\pm$ 1.30 <sup>a</sup>
Extract 400 mg	2.54 $\pm$ 0.95 <sup>a</sup>	4.34 $\pm$ 1.74 <sup>a</sup>	9.2 $\pm$ 3.42 <sup>b</sup>

Note: Results represent mean  $\pm$  standard deviation of group results (n=5). Values having different superscripts along the column are considered statistically significant ( $p < 0.05$ )



**Table 4.** Effect of *Phyllanthus amarus* leaves extraction on some liver function biomarkers of wistar rat

Treatment	ALB (mg/mL)	TP (mg/mL)	DB ( $\mu\text{mol/L}$ )	TB ( $\mu\text{mol/L}$ )
Control	2.6 $\pm$ 1.25 <sup>a</sup>	8.98 $\pm$ 3.07 <sup>a</sup>	30.03 $\pm$ 6.34 <sup>ab</sup>	30.96 $\pm$ 7.70 <sup>a</sup>
Extract 100 mg	2.78 $\pm$ 0.38 <sup>a</sup>	8.74 $\pm$ 1.21 <sup>a</sup>	30.36 $\pm$ 4.58 <sup>ab</sup>	34.66 $\pm$ 9.89 <sup>a</sup>
Extract 200 mg	3.2 $\pm$ 0.72 <sup>a</sup>	9.62 $\pm$ 2.82 <sup>a</sup>	33.56 $\pm$ 9.11 <sup>b</sup>	36.84 $\pm$ 6.50 <sup>a</sup>
Extract 400 mg	3.42 $\pm$ 0.61 <sup>a</sup>	11.54 $\pm$ 5.82 <sup>a</sup>	23.84 $\pm$ 4.33 <sup>a</sup>	56.86 $\pm$ 17.63 <sup>a</sup>

Note: Results represent mean  $\pm$  standard deviation of group results (n=5). Values having different superscripts along the column are considered statistically significant ( $p < 0.05$ )

## Discussion

Kidneys and the liver are vital organs that function for homeostasis, regulating electrolyte balance in the blood, excreting waste products of metabolism, secretion of some enzymes and hormones, metabolism, osmoregulation, and detoxification. However, damage and toxic substances in these essential organs can lead to inefficiency in carrying out their functions.

Electrolytes (sodium, potassium, chloride, and bicarbonate) balance in the blood is a good indicator of kidneys and heart functions. Therefore, adequate information on serum electrolytes can assist in the determination of organ functions. For example, liver function assessment can be done by evaluating liver enzyme (AST, ALT, and ALP) parameters, total protein, albumin, and total bilirubin (Yakubu et al. 2021a).

The result revealed a significant decrease ( $p > 0.05$ ) in sodium ion ( $\text{Na}^+$ ) concentration between groups administered with doses of extract (100 mg, 200 mg) compared to normal/ control. Still, it showed no significant ( $p > 0.05$ ) difference at 400 mg dosage compared to the normal/ control. Therefore, the results suggested mild to no change of sodium ion concentration in the serum of rats administered with *P. amarus*. Furthermore, the result also revealed no significant ( $p > 0.05$ ) change in the concentration level of the other electrolytes, i.e., chlorine ( $\text{Cl}^-$ ) and Potassium ( $\text{K}^+$ ), between animal groups administered with the extract and normal control. These findings imply that the administration of *P. amarus* extract within these experimental doses did not significantly change the electrolyte concentrations in the serum of the wistar rats. This result agrees with the work of Krithika and Verma (2009) that the LD<sub>50</sub> value of *P. amarus* was high or categorized as non-toxic. The result also revealed no significant difference in serum urea and creatinine concentration between and within the groups of the administered dose of 100 mg, 200 mg, and 400 mg compared to normal control. Hence, the serum's electrolytes values, urea, and creatinine data indicate that the extract did not exact any deleterious effect on the kidney or negatively affect the heart.

The liver is an essential organ that carries out the biological system's metabolic, secretion, storage, and detoxification functions. Therefore, direct or indirect damage to the hepatocytes will cause an alteration of these functions, thereby affecting the organism's living condition (Yakubu et al. 2020b). The liver parameters showed no significant ( $p > 0.05$ ) difference in the level of ALB, TP, TB, DB, ALT, AST, and ALP of the animal groups

administered with extract compared to the normal control. Based on the parameters observed, it shows that the administration of the extract did not cause damage or toxicity to the liver. *P. amarus* contains alkaloids and a high concentration of antioxidant compounds (Fernand 1998; Naaz et al. 2007; Krithika and Verma 2009), essential for reducing oxidative stress. It may also have been the major factor in stabilizing all the hepatic parameters (Yakubu et al. 2021b). However, inappropriate use of antioxidants can induce oxidative stress (Galati and O'Brien 2004; Atici et al. 2005); hence the use of *P. amarus* extract should be kept at an optimal level.

It can be concluded that administration of *P. amarus* extract up to 400 mg/kg BW did not induce hepatotoxicity or nephrotoxicity. Furthermore, this finding has validated the use of *P. amarus* as a traditional herb for treating and managing several diseases, including kidney and liver diseases, without any significant organ damage.

## REFERENCES

- Amador E, Wacker W. 1962. Analytical methods for quantitative determination of liver marker enzymes. Clin Chem 8: 343. DOI: 10.1093/clinchem/8.4.343.
- Atici S, Cinel I, Cinel L, Doruk N, Eskandari G, Oral U. 2005. Liver and kidney toxicity in chronic use of opioids: An experimental long term treatment model. J Biosci 30: 245-252. DOI: 10.1007/BF02703705.
- Burnett RW, Covington AK, Fogh-Andersen N, Kùlpmann WR, Lewenstam A, Maas AH et al. 2000. Recommendations for measurement of and conventions for reporting sodium and potassium by ion-selective electrodes in undiluted serum, plasma or whole blood. International Federation of Clinical Chemistry and Laboratory Medicine (IFCC). IFCC Scientific Division Working Group on Selective Electrodes. Clin Chem Lab Med 38: 1065-1071. DOI: 10.1515/CCLM.2000.159.
- Calixto JB, Santos ARS, Cechinel-Filho V, Yunes RA. 1998. A review of the plants of the Genus *Phyllanthus*: Their chemistry, pharmacology and therapeutic potential. Med Res Rev 18: 225-258. DOI: 10.1002/(sici)1098-1128(199807)18:4<225::aid-med2>3.0.co;2-x.
- Emeka O, Olotu EJ, Allison TA, Onyekwere JC. 2013. The wound healing effects of aqueous leave extracts of *Azadirachta Indica* on wistar rats. J Nat Sci Res 3 (6): 181-186.
- Fernand VE. 1998. Initial Characterization of Crude Extracts from *Phyllanthus amarus* Schum and Thonn and *Quassia Amara* L. sing Normal Phase Thin Layer Chromatography. [Thesis]. Louisiana State University, United States of America.
- Galati G, O'Brien PJ. 2004. Potential toxicity of flavonoids and other dietary phenolics: Significance for their chemopreventive and anticancer properties. Free Radic Biol Med 37: 287-303. DOI: 10.1016/j.freeradbiomed.2004.04.034.
- Haussament TU. 1977. Quantitative determination of serum alkaline phosphatase. Clinica Chimica Acta 35: 271-273.
- Ilonga SK, Kandawa-Schulz M, El-Sayed LoftyH, Lyantagaye S. 2018. Anticancer, antioxidant and antimicrobial screening of extracts from

- selected medicinal plants from Oshikoto, Namibia. *Bioteknologi* 15: 51-65. DOI: 10.13057/biofar/c150202.
- Krithika R, Verma RJ. 2009. Mitigation of carbon tetrachloride-induced damage by *Phyllanthus amarus* in liver of mice. *Acta Poloniae Pharm* 66 (4): 439-444.
- Liu RH. 2003. Health benefits of fruit and vegetables are from additive and synergistic combinations of phytochemicals. *Am J Clin Nutr* 78: 517S-520S. DOI: 10.1093/ajcn/78.3.517S.
- Matthews HB, Lucier GW, Fisher KD. 1999. Medicinal herbs in the United States: Research Needs. *Environ Health Perspect* 107: 773-778. DOI: 10.1289/ehp.99107773.
- Naaz F, Javed S, Abdin MZ. 2007. Hepatoprotective effect of ethanolic extract of *Phyllanthus amarus* Schum. et Thonn. on aflatoxin B<sub>1</sub>-induced liver damage in mice. *J Ethnopharmacol* 113: 503-509. DOI: 10.1016/j.jep.2007.07.017.
- Srividiya N, Perival S. 1995. Diuretic, hypotensive and hypoglycemic effect of *Phyllanthus amarus*. *Indian J Exp Biol* 33 (11): 861-864.
- Tuhinadri S, Sami KS. 2015. Medicinal plants, human health and biodiversity: A broad review. *Adv Biochem Eng Biotechnol* 147: 59-110. DOI: 10.1007/10\_2014\_273.
- Yakubu OE, Imo C, Shaibu C, Akighir J, Ameh DS, 2020a. Effects of ethanolic leaf and stem-bark extracts of *Adansonia digitata* in alloxan-induced diabetic wistar rats. *J Pharmacol Toxicol* 15: 1-7. DOI: 10.3923/jpt.2020.1.7.
- Yakubu OE, Nwode OFC, Joshua PE, Ugwu MN, Odu AD, Okwo F. 2014. Fractionation and determination of total antioxidant capacity, total phenolics and flavonoids content of aqueous, ethanol and n-hexane extract of *Vitexdonia leave*. *Afr J Biotechnol* 13 (5): 693-698. DOI: 10.5897/AJB2013.13225.
- Yakubu OE, Ojogbane E, Nwaneri-Chidozie VO, Ibuzo PN, Ale EM, Awudu SS. 2020b. Ameliorative effects of ethanol extract of ficus sur on diethylnitrosamine-induced hepatotoxicity in wistar rats. *Am J Biochem Mol Bio* 10: 6-11. DOI: 10.3923/ajbmb.2020.6.11.
- Yakubu OE, Ojogbane EB, Abu MS, Shaibu CO, Ayegba WE. 2020c. Haematinic effects of ethanol extract of *Ficus sur* leaves on diethylnitrosamine-induced toxicity in wistar rats. *J Pharm Pharmacol* 15 (1): 16-21. DOI: 10.3923/jpt.2020.16.21.
- Yakubu OE, Ojogbane EB, Atanu FO, Udeh CSM, Ale ME, Bello BH. 2021a. Hepatoprotective effects of partially purified fractions of *Senna occidentalis* ethanolic extract on diethyl nitrosamine-induced toxicity in wistar rats. *Intl J Sci Rep* 7 (8): 374-81. DOI: 10.18203/issn.2454-2156.IntJSciRep20212837.
- Yakubu OE, Udeh CS, Asuk AA, Moses AA, Richard B. 2021b. Comparative determination of antioxidant activities and phytochemicals from fractions of ethanol extract of *Senna occidentalis* using GC-MS. *J Emerg Technol Innov Res* 8 (7): 195-211.

## Comparison of phytochemical constituents of ethanol leaf extracts of *Solanum macrocarpon* and *Vernonia amygdalina*

OLUSOJI ADEBUSOYE OYESOLA<sup>1</sup>, IQOT ISAAC SAMPSON<sup>1</sup>, ADEBIYI ADELOWO AUGUSTINE<sup>2</sup>,  
OLUKADE BALIQIS ADEJOKE<sup>1</sup>, GEORGE EMMANUEL TAIWO<sup>1,\*</sup>

<sup>1</sup>Department of Physiology, Faculty of Basic Medical Sciences, Olabisi Onabanjo University, Ago Iwoye, Ogun State, Nigeria.  
Tel./fax. +234-805-560 6475, \*email: georgeayoku@gmail.com

<sup>2</sup>Department of Biochemistry, Faculty of Basic Medical Sciences, Olabisi Onabanjo University, Ago Iwoye, Ogun State, Nigeria

Manuscript received: 18 February 2022. Revision accepted: 9 April 2022.

**Abstract.** Oyesola OA, Sampson II, Augustine AA, Adejoke OB, Taiwo GE. 2022. Comparison of phytochemical constituents of ethanol leaf extracts of *Solanum macrocarpon* and *Vernonia amygdalina*. *Asian J Trop Biotechnol* 20: 6-10. A plant can exert physiological changes on biological systems due to phytochemicals present in plant parts. This study evaluated and compared the phytochemical constituent (both quantitative and qualitative) and the antioxidant properties of the ethanol leaf extract *Solanum macrocarpon* L. and *Vernonia amygdalina* Delile using standard methods. Tannin, phenol, cardiac glycoside, alkaloid, and flavonoid were present in both plant extracts. In large quantities of alkaloid, flavonoid, and cardiac glycoside in *S. macrocarpon* compared to *V. amygdalina*. At the same time, *V. amygdalina* contained a large amount of tannin and phenol. Reducing sugar was present in *S. macrocarpon* and absent in *V. amygdalina*; saponin and steroid were present in *V. amygdalina* and absent in *S. macrocarpon*; terpenoid and phlobatanin were both missing in the leaf extract of both plants. In contrast, both plant extracts showed radical scavenging activities, with *V. amygdalina* having a higher antioxidant capacity than *S. macrocarpon*. The present study's findings indicated that the ethanol leaf extract of *S. macrocarpon* and *V. amygdalina* possess antioxidant properties and may be effective against oxidative stress.

**Keywords:** Antioxidant, phytochemical, *S. macrocarpon*, *V. amygdalina*

### INTRODUCTION

Plants' medicinal properties are attributed to synthetic components that have a distinct physiological effect on the human body; these compound substances are known as phytochemicals (Edeoga et al. 2005). Phytochemicals are currently receiving more attention because of their effectiveness in treating infectious diseases (Wamuyu et al., 2020). The phytochemical analysis of medicinal plants involves extracting, screening, and identifying bioactive compounds in different plant parts. Flavonoids, alkaloids, carotenoids, tannins, antioxidants, and phenolic compounds are some bioactive molecules obtained from plants. Phytochemicals are naturally found in plants and play an essential role in assisting plants to protect themselves against pathogenic microbes by demonstrating antimicrobial activity through hindrance or killing mechanisms. The discharge of these mixtures varies from one plant to the next, with some producing higher quality and others producing low quality (Tariq and Reyaz 2013). Research has found that phytochemicals present in fruits and vegetables decrease the risk of cancer, act as antioxidants, and may treat or manage infections and metabolic disorders (Abbasi et al., 2015). *Solanum macrocarpon* L. is the scientific name for the African eggplant, which belongs to the Solanaceae family and the plant genus *Solanum* (Agoreyo et al. 2012). Eggplants have a wide range of nutritional and therapeutic properties, making them a beneficial complement to any diet because

they contain a significant amount of nutrients and phytochemical compounds such as saponins, phenols, flavonoids, and tannins, among others (Ibiam and Nwigwe 2013). Eggplant fruit aids in the prevention, management, and treatment of various diseases by lowering blood cholesterol levels, managing high blood pressure, reducing weight, and having anti-haemorrhoidal and anti-glaucoma properties (Ossamulu et al. 2014).

Because of its bitter taste, *Vernonia amygdalina* Delile is also called a bitter leaf. A small evergreen shrub grows throughout Africa and belongs to the Asteraceae family. It was reported to be a plant that can help with diabetes and fever management (Imaga and Bamigbetan 2013). The bitter taste of *V. amygdalina* is due to the presence of sesquiterpene lactones (vernodaline, vernolepin, and vernomygdin) and steroid glycosides (vernonisides) (Ojmelukwe and Amaechi 2019). Previous phytochemical studies on the leaf extract of *V. amygdalina* shows the presence of bioactive compounds like tannins, saponins, flavonoid, glycosides, alkaloids, and steroid. Traditionally, plants are also used to treat and manage malaria, intestinal parasite, diarrhea, and high blood sugar (Udochukwu et al., 2015). African eggplant and bitter leaf are important vegetables in African communities due to their nutritional and medicinal value; the study will compare and analyze the phytochemical constituent and in-vitro antioxidant properties of the ethanol leaf extract of *S. macrocarpon* and *V. amygdalina*; it will also add to the knowledge on their significance use in ethnomedicine.

## MATERIALS AND METHODS

### Plants material

Matured leaves of bitter leaf plants were collected from a family garden in the Sagamu local government area of South-West Nigeria. In contrast, matured leaves of African eggplant were bought from a local market in the Sagamu local government area of South-West Nigeria. In addition, authentications of the leaves sample were carried out at the department of plant science, faculty of science, Olabisi Onabanjo University, Nigeria.

### Preparation of the ethanol leaves extract of African eggplant and bitter leaf plant

The leaves of African eggplant and bitter leaf plant were air-dried and powdered using a blender; 150 g of the blended leaves were soaked in 750 mL of ethanol (70% ethanol and 30% water) for three days and then filtered. After filtration, the filtrate was heated at a temperature of 40°C for evaporation.

### Determination of percentage yield of African eggplant and bitter leaf plant

The percentage of African eggplant extract was determined by calculating the percentage of the weight of the extract to the original weight before drying the sample, using the formula;

$$\text{percentage yield} = \frac{\text{weight of extract}}{\text{weight of sample}} \times \frac{100}{1}$$

Weight of African eggplant= 150g

Weight of dried shaft of African eggplant= 73.4g

Weight of extract = 150g — 73.4g= 76.6g

$$\text{percentage yield} = \frac{76.6\text{g}}{150\text{g}} \times \frac{100}{1} = 51.0\%$$

The percentage yield for African eggplant is 51.0%

The percentage yield for bitter leaf plant was also calculated using the same formula stated above;

$$\text{percentage yield} = \frac{\text{weight of extract}}{\text{weight of sample}} \times \frac{100}{1}$$

Weight of bitter leaf plant – 150g

Weight of dried shaft of bitter leaf plant- 70.10g

Weight of extract = 150g—70.10g= 79.9g

$$\text{percentage yield} = \frac{79.9\text{g}}{150\text{g}} \times \frac{100}{1} = 53.3\%$$

The percentage yield for bitter leaf is 53.3g

### Phytochemical screening and in-vitro antioxidant procedure

Phytochemical tests were carried out on the ethanol leaf extract of *V. amygdalina* and *S. macrocarpon* using the standard procedure to identify the constituents present and

in-vitro antioxidant enzymes activity as described by Harborne (1973), Trease and Evans (1989), Sofowra (1993), and Alisi and Onyeze (2008).

## RESULTS AND DISCUSSION

### Determination of the phytochemical present in the ethanol extract of the leaves extract of *Solanum macrocarpon* and *Vernonia amygdalina*

Tables 1 and 2 show the qualitative and quantitative (mg/100 g) analysis of the phytochemical constituents in the ethanol leaf extract of *S. macrocarpon* and *V. amygdalina*. The results revealed the presence of bioactive compounds in the extract studied from the table; the results show that phenols and tannins, flavonoids, and cardiac glycoside were present in both plant extracts. However, saponins were absent only in the ethanol extract of the African eggplant, and reducing sugar was absent in the ethanol extract of the bitter leaf plant. At the same time, steroids were missing in the ethanol extract of African eggplant. In addition, terpenoid and philobatanin were not present in both plant extracts. The results also show that the leaves of bitter leaf contained high levels of tannin and phenol compared to the leaves of African eggplant. In contrast, the leaves of African eggplant contain more alkaloids, cardiac glycoside, and flavonoids compared with the leaves of bitter leaf.

Plant phytochemical constituents are increasingly linked to the elicited physiological activities; in traditional medicine, plant parts are used to manage and treat various disorders (Gurib-Fakim 2006). The phytochemical screening results in Table 3 show the presence of active entities that elicit significant pharmacological and physiological responses. The presence of alkaloids, flavonoids, phenolic compounds, tannins, saponins, terpenoids, reducing sugar, steroid, and cardiac glycosides was observed. Tannins and flavonoids in the plant extract are responsible for the observed DPPH radical scavenging activity, as seen in Table 3. Flavonoids and tannins are phenolic compounds, which are the most abundant bioactive compounds in plants that act as antioxidants or free radical scavengers. The presence of one bioactive compound in the plant extract and its absence in another may be due to a difference in solvent polarity, which follows the rules of thumb" like dissolves like" (Adamu et al. 2019). The bitterness of African eggplants and bitter leaf plants is caused by the presence of alkaloids, primarily glycoalkaloids, and the degree of bitterness determines the edibility or otherwise. In general, ethanol extracts of bitter leaf plants contained more phytochemicals than ethanol extracts of African eggplant. The saponins found in the samples are major nutritional substances and nutraceuticals. Previous research studies have shown that medicinal plants' saponins reduce glycoside toxicity by hydrolyzing terpenoids (Xu et al. 1996; Chinedu et al. 2011). The antioxidant activity shown by the plant extracts in Tables 3-6 may result from the presence of phenolic compounds. The antioxidant action of phenolic compounds stems from their redox characteristics, which can aid the

absorption and neutralization of free radicals, singlet and triplet oxygen quenching, and the decomposition of peroxides (Al-Shaya et al., 2020). The food industry is increasingly interested in crude extracts of high phenolic medicinal plant materials (Osawa 1994).

Alkaloids are essential for plant protection and survival because they protect them from microorganisms' activities, insects and herbivores, and other plants (allelopathically active chemicals) (Molyneux et al. 1996). Plants containing alkaloids have been used as dyes, spices, drugs, and poisons almost since the beginning of human history. Cardiac glycosides are derived from steroids and act primarily on the cardiac muscle; they are potent in

managing heart disease. Congestive heart failure causes an influx of Na<sup>+</sup> and an outflow of K<sup>+</sup> during each heart contraction. Na<sup>+</sup>, K<sup>+</sup>-ATPase must re-establish the concentration gradient before the next contraction by pumping Na<sup>+</sup> into the cell against a concentration gradient. Cardiac glycosides inhibit Na<sup>+</sup> K<sup>+</sup>-ATPase, increasing and increasing the force of myocardial contraction as a result (Farnsworth 1966); cardiac glycosides also have antitumor activity (Dorskotch et al. 1972). Other studies have also reported the presence of a physiologically active substance in bitter leaf (Usunobun and Okolie 2015; Usunomena and Ngozi 2016) and African eggplant (Iiodibia et al. 2016; Eletta et al. 2017).

**Table 1.** Quantitative determination of phytochemical constituents of ethanol leaf extract African eggplant and bitter leaf plant

Sample	Tannin mg/100 g	Alkaloid mg/100 g	Reducing sugar mg/100 g	Cardiac glycoside mg/100 g	Phenol mg/100 g	Steroid mg/100 g	Flavanoid mg/100 g	Saponin mg/100 g
<i>V. amygdalina</i>	45.09	39.14	-	38.11	61.93	-	51.93	51.99
<i>S. macrocarpon</i>	39.80	69.61	45.90	38.30	41.21	31.25	67.51	-

**Table 2.** Qualitative determination of phytochemical constituents of ethanol leaf extracts African eggplant and bitter leaf plant

Sample	Tannin	Alkaloid	Reducing sugar	Cardiac glycoside	Terpenoid	Phenol	Phlobatanin	Steroid	Flavanoid	Saponin
<i>V. amygdalina</i>	+	+	-	+	-	+	-	+	+	+
<i>S. macrocarpon</i>	+	+	+	+	-	+	-	-	+	-

Note: + =present, - =absent

**Table 3.** DPPH radical scavenging activity of ethanol leaf extracts of *Solanum macrocarpon* and *Vernonia amygdalina*

Sample	25 µg/mL	50 µg/mL	75 µg/mL	100 µg/mL
<i>V. amygdalina</i>	31.25	47.06	59.44	77.72
<i>S. macrocarpon</i>	33.65	50.22	56.90	63.50

**Table 4.** Nitric oxide scavenging activity of ethanol leaf extracts *Solanum macrocarpon* and *Vernonia amygdalina*

Sample	25 µg/mL	50 µg/mL	75 µg/mL	100 µg/mL
<i>V. amygdalina</i>	37.88	48.07	61.29	77.27
<i>S. macrocarpon</i>	38.22	41.89	53.06	61.66

**Table 5.** Reducing power of ethanol leaf extracts *Solanum macrocarpon* and *Vernonia amygdalina*

Sample	25 µg/mL	50 µg/mL	75 µg/mL	100 µg/mL
<i>V. amygdalina</i>	0.12	0.26	0.32	0.46
<i>S. macrocarpon</i>	0.17	0.21	0.36	0.48

**Table 6.** Total flavonoid, total phenol, and total antioxidant activity of ethanol leaf extract *Solanum macrocarpon* and *Vernonia amygdalina*

Sample	Total flavonoid mg/100 g	Total phenol Mg/100 g	Total antioxidant capacity
<i>V. amygdalina</i>	51.92	61.93	52.07
<i>S. macrocarpon</i>	67.51	27.45	39.71

### Evaluation of the antioxidant activity of ethanol leaves extract of *Solanum macrocarpon* and *Vernonia amygdalina*

Total Antioxidant Capability (TAC) was coined to describe the antioxidant's reducing capacity in a single metric. TAC of the ethanol extracts of *S. macrocarpon* and *V. amygdalina* calculated using a variety of methodologies, including a DPPH radical scavenging activity assay and a nitric oxide scavenging activity, and a reducing power assay. In the DPPH radical scavenging activity assay, nitric oxide scavenging activity, and reducing power assay, absorbance directly represents reducing power (McCord 2000). Tables 3-6 show the ethanol extract's antioxidant activities in bitter leaf and African eggplant. The results of the DPPH scavenging activity of the extracts are shown in Table 4. The African eggplant and bitter leaf plant ethanol extract exhibited concentration-dependent antiradical activity by inhibiting DPPH radical with inhibitory concentrations of 63.50 at 100g/ml and 77.72 at 100 g/mL, respectively. Researchers commonly utilize the model system of DPPH radicals to explore the scavenging ability of various medicinal plant products (Benslama and Harrar 2016). Due to the hydrogen-donating ability of antioxidants, they can scavenge DPPH radicals (Baumann 1979). Free radical scavengers are important to prevent the harmful effects of free radicals. The DPPH free radical scavenging methods are widely used to evaluate the antioxidant properties of plants extracts; the DPPH radical scavenging activity of *V. amygdalina* corresponds with the study of Erasto et al. (2007), Ho et al. (2012), Atangwho et al. (2013), while that of *S. macrocarpon* correspond with the study of Adewale et al. (2014) and Eletta et al. (2017). In the reducing power assay (Table 5), the presence of antioxidants in the samples would result in the extract donating an electron to reduce  $Fe^{3+}$  to  $Fe^{2+}$ . The extracts with reducing power reveal that they are electron donors, reduce oxidized intermediates, and act as primary antioxidants (Chanda and Dave 2009). The reducing power is frequently employed to access the antioxidant activity of natural plant products. The presence of reductants which act as an antioxidant by breaking free radical chains by donating a hydrogen atom, is often associated with the existence of reducing power (Rahman et al., 2015). Observation from Table 5 shows that the reducing power of both plant extracts was in different concentrations. These results also correspond with other studies which show that *S. macrocarpon* (Adewale et al. 2014; Famuwagun et al. 2017) and *V. amygdalina* (Ho et al. 2012; Adesanoye and Farombi 2014) have reduced power activity, but the ethanol leaf extract of *S. macrocarpon* had more reducing power when compared to the ethanol leaf extract of *V. amygdalina*. Nitric oxide (NO) or reactive nitrogen species such as  $NO_2$ ,  $N_2O_4$ ,  $N_3O_4$ ,  $NO_3$ , and  $NO_2$  are formed during reactive nitrogen reactions with oxygen or superoxides. These compounds alter many cellular components' structural and functional behavior. Plant products can inhibit the detrimental consequences of excessive NO production in the human body, which could be of considerable interest in preventing the harmful effects of excessive NO production. NO has

also been linked to inflammation, cancer, and other diseases (Moncada and Higgs 1993). Table 5 shows the relative NO scavenging potential of African eggplant and bitter leaf plant ethanol extracts. Since its identification as a new signal molecule, NO is linked to various physiological reactions. It produces vascular dilation by transmitting signals from vascular endothelial cells to vascular smooth muscle cells. It also plays an important part in respiratory, immunological, neuromuscular, and other physiological activities (Ebrahimzadeh et al., 2010). Other studies reported the NO scavenging activity of *S. macrocarpon* and *V. amygdalina* (Ng et al., 2015; Omede et al., 2018). Phenols are secondary metabolites found in plants. Most plant products can cause a pharmacological effect which includes; anti-inflammatory, antispasmodic, and anti-allergic, among others. Most pathological diseases and infections, such as diabetes, cancer, and cardiovascular disease resulting from oxidative injury, are caused by oxidative stress. The accumulation of reactive oxygen species (ROS) triggers a series of reactions that breaks down organic molecule such as DNA, lipids, and protein in the body, which are the main causes of disease (Halliwell et al. 1992; Craig 1999; Exarchou et al. 2002; Afanas'ev 2010). Many plant extracts that contain phenol compounds possess antimicrobial and antioxidant properties and are used to treat and manage disease (Choi et al., 2010). In Table 6, it was observed that the ethanol leaf extract of *V. amygdalina* has a high quantity of phenols compared with the ethanol leaf extract of *S. macrocarpon*. At the same time, total flavonoid content was high in the ethanol leaf extract of *S. macrocarpon* compared with *V. amygdalina*. The antioxidant activities of these plant extracts are due to the presence of flavonoid and phenol compounds.

In conclusion, the presence of tannins, alkaloids, cardiac glycoside, phenol, and flavanoid in the ethanol leaf extracts of *V. amygdalina* and *S. macrocarpon* are responsible for the physiological changes exerted by the plant on biological systems and also possess antioxidant properties and may be effective against oxidative stress caused by free radicals. However, more studies are needed to isolate the active ingredients in the combination of African eggplant and bitter leaf ethanol extracts and study their anti-inflammatory effect.

### REFERENCES

- Abbasi T, Anuradha J, Ganaie SU, Abbasi SA. 2015. Biomimetic synthesis of nanoparticles using aqueous extracts of plants (Botanical Species). *J Nano Res* 31: 138-202. DOI: 10.4028/www.scientific.net/JNanoR.31.
- Adamu HM, Yushau S, Yakubu H, Abubakar A. 2019. Phytochemical screening and antioxidant activity of the stem bark extracts of *Diospyros mespiliformis*: A medicinal plant in Bauchi. *Intl J Chem Sci* 4 (3): 37-42. DOI: 10.31838/ijprt/10.01.08.
- Adesanoye OA, Farombi EO. 2014. In vitro antioxidant properties of methanolic leaf extract of *Vernonia amygdalina* Del. *Niger J Physiol Sci* 29 (2): 93-101.
- Adewale OB, Onasanya A, Fadaka AO, Iwere H, Anadozie SO, Osukoya OA, Olayide II. 2014. In vitro antioxidant effect of aqueous extract of *Solanum macrocarpon* leaves in rat liver and brain. *Oxid Antioxid Med Sci* 3 (3): 225-229. DOI 10.5455/oams.191214.or.079.
- Afanas'ev I. 2010. Signaling and damaging functions of free radicals in aging-free radical theory, hormesis, and TOR. *Aging Dis* 1 (2): 75-88.

- Agoreyo BO, Obansa ES, Obanor EO. 2012. Comparative nutritional and phytochemical analyses of two varieties of *Solanum melongena*. *Sci World J* 7 (1): 5-8. DOI: 10.4314/bajopas.v5i1.32.
- Alisi CS, Onyeze GOC. 2008. Nitric oxide scavenging ability of ethyl acetate fraction of methanolic leaf extracts of *Chromolaena Odorata* (Linn.). *Afr J Biochem Res* 2 (7):145-150. DOI: 10.5897/AJBR.9000174.
- Al-Shaya HM, Li H, Beg OU, Hamama AA, Witiak SM, Kaseloo P, Siddiqui RA. 2020. Phytochemical profile and antioxidation activity of *Annona* fruit and its effect on lymphoma cell proliferation. *Food Sci Nutr* 8 (1): 58-68. DOI: 10.1002/fsn3.1228.
- Atangwho II, Egbung GE, Ahmad M, Yam MF, Asmawi MZ. 2013. Antioxidant versus anti-diabetic properties of leaves from *Vernonia amygdalina* Del. growing in Malaysia. *Food Chem* 141 (4): 3428-3434. DOI: 10.1016/j.foodchem.2013.06.047.
- Baumann J. 1979. Prostaglandin synthetase inhibiting O<sub>2</sub>-radical scavenging properties of some flavonoids and related phenolic compounds. *Naunyn-Schmiedeberg Arch Pharmacol* 308: 27-32.
- Benslama A, Harrar A. 2016. Free radicals scavenging activity and reducing power of two Algerian Sahara medicinal plants extracts. *Intl J Herb Med* 4 (6): 158-161. DOI: 10.22271/flora.2016.v4.i6c.03.
- Chanda S, Dave R. 2009. In vitro models for antioxidant activity evaluation and some medicinal plants possessing antioxidant properties: An overview. *Afr J Microbiol Res* 3 (13): 981-996. DOI: 10.5897/AJMR.9000401.
- Chinedu SN, Olasumbo AC, Eboji OK, Emiloju OC, Arinola OK, Dania DT. 2011. Proximate and phytochemical analyses of *Solanum aethiopicum* L. and *Solanum macrocarpon* L. fruits. *Res J Chem Sci* 1(3): 63-71.
- Choi BS, Sapkota K, Kim S, Lee HJ, Choi HS, Kim SJ. 2010. Antioxidant activity and protective effects of *Tripterygium regelii* extract on hydrogen peroxide induced injury in human dopaminergic cells, SH-SY5Y. *Neurochem Res* 35: 1269-1280. DOI: 10.1007/s11064-010-0185-4.
- Craig WJ. 1999. Health-promoting properties of common herbs. *The American journal of clinical nutrition*. 70 (3): 491s-499s. DOI: 10.1093/ajcn/70.3.491s.
- Doskotch RW, Malik MY, Hufford CD, Malik SN, Trent JE, Kubelka W. 1972. Antitumor agents. V: Cytotoxic cardenolides from *Cryptostegia grandiflora* (Roxb.) R. Br. *J Pharm Sci* 61: 570-573. DOI: 10.1002/jps.2600610415.
- Ebrahimzadeh MA, Nabavi SF, Nabavi SM, Pourmorad F. 2010. Nitric oxide radical scavenging potential of some Elburz medicinal plants. *Afr J Biotechnol* 9 (32): 5212-7.
- Edeoga HO, Okwu DE, Mbaebie BO. 2005. Phytochemical constituents of some Nigerian medicinal plants. *Afr J Biotechnol* 4 (7): 685-688. DOI: 10.5897/AJB2005.000-3127.
- Eletta OAA, Orimolade BO, Oluwaniyi OO, Dosumu OO. 2017. Evaluation of proximate and antioxidant activities of Ethiopian eggplant (*Solanum aethiopicum* L) and Gboma eggplant (*Solanum macrocarpon* L). *J Appl Sci Environ Manag* 21 (5): 967-972. DOI: 10.4314/jasem.v21i5.25.
- Erasto P, Grierson DS, Afolayan AJ. 2007. Evaluation of antioxidant activity and the fatty acid profile of the leaves of *Vernonia amygdalina* growing in South Africa. *Food Chem* 104 (2): 636-642. DOI: 10.1016/j.foodchem.2006.12.013.
- Exarchou V, Nenadis M, Tsimidou IP, Gerothanassis A, Troganis D, Boskou. 2002. Antioxidant activities and phenolic composition of extracts from Greek oregano, Greek sage, and Summer savory. *J Agric Food Chem* 50: 5294-5299. DOI: 10.1021/jf020408a.
- Famuwagun AA, Taiwo KA, Gbadamosi SO, Oyedele DJ, Aluko RE, Adebooye OC. 2017. Extraction optimization and antioxidant properties of African eggplant (*Solanum macrocarpon*) leaf polyphenols. *J Food Qual* 2017 (3): 1-14. DOI: 10.1155/2017/2159183.
- Farnsworth NF. 1966. Biological and phytochemical screening of plants. *J Pharm Sci* 55: 225-276. DOI: 10.1002/jps.2600550302.
- Gurib-Fakim A. 2006. Medicinal plants: Traditions of yesterday and drugs of tomorrow. *Mol Asp Med* 27 (1): 1-93. DOI: 10.1016/j.mam.2005.07.008.
- Halliwell B, Gutteridge JMC, Cross CE. 1992. Free radicals, antioxidants, and human disease: Where are we now?. *J Lab Clin Med* 119: 598-620.
- Harborne JB. 1973. *Phytochemicals Methods*. Chapman and Hall Ltd., London.
- Ho WY, Liang WS, Yeap SK, Beh BK, Youns AH, Alitheen NB. 2012. In vitro and in vivo antioxidant activity of *Vernonia amygdalina* water extract. *Afr J Biotechnol* 11 (17): 4090-4094. DOI: 10.5897/AJB10.1639.
- Ibiam, OF, Nwigwe I. 2013. The effect of fungi associated with leaf blight of *Solanum aethiopicum* L. in the field on the nutrient and phytochemical composition of the leaves and fruits of the plant. *J Plant Pathol Microbiol* 4 (7): 191-195. DOI: 10.4172/2157-7471.1000191.
- Iloobia CV, Akachukwu EE, Chukwuma MU, Igboabuchi NA, Adimonyemma RN, Okeke, NF. 2016. Proximate, phytochemical and antimicrobial studies on *Solanum macrocarpon* L. *J Adv Biol Biotechnol* 9 (2): 1-7. DOI: 10.9734/JABB/2016/27922.
- Imaga NOA, Bamigbetan DO. 2013. In vivo biochemical assessment of aqueous extracts of *Vernonia amygdalina* (Bitter leaf). *Intl J Nutr Metabol* 5 (2): 22-27. DOI: 10.5897/IJNAM12.0001.
- McCord JM. 2000. The evolution of free radicals and oxidative stress. *J Med* 108: 652-659. DOI: 10.1016/s0002-9343(00)00412-5.
- Molyneux RJ, Nash RJ, Asano N. 1996. The chemistry and biochemistry of simple indolizidine and related polyhydroxy alkaloids. In: Pelletier SW (eds). *Alkaloids: Chemical and Biological Perspectives*, Vol. 11. Pergamon, Oxford.
- Moncada S, Higgs A. 1993. Mechanisms of diseases: The largininenitricoxide pathway. *N Engl J Med* 329 (27): 2002-2012. DOI: 10.1056/NEJM199312303292706.
- Ng RF, Abidin NZ, Shuib AS, Israf Ali DA. 2015. Inhibition of nitric oxide production by *Solanum melongena* and *Solanum macrocarpon* on RAW 264.7 cells. *Front Life Sci* 8 (3):241-8. DOI: 10.1080/21553769.2015.1051241.
- Ojimelukwe PC, Amaechi N. 2019. Composition of *Vernonia amygdalina* and its potential health benefits. *Intl J Environ Agric Biotech* 4 (6): 1836-1848. DOI: 10.22161/ijeab.46.34.
- Omede A, Suleiman MS, Atanu FO, Momoh S, Friday ET, Sheneni VD, Jegede ER. 2018. Evaluation of antioxidant and cytotoxic properties of *Vernonia amygdalina*. *Intl J Cell Sci Mol Biol* 4 (4): 81-86. DOI: 10.19080/IJCSMB.2018.03.555644.
- Osawa T. 1994. Novel natural antioxidants for utilization in food and biological systems. In: Utritani I, Garcia VV, Mendoza EM. (eds). *Postharvest Biochemistry of Plant Food Materials in the Tropics*. Japan Scientific Societies Press, Tokyo.
- Rahman MM, Islam MB, Biswas M, Alam AK. 2015. In vitro antioxidant and free radical scavenging activity of different parts of *Tabebuia pallida* growing in Bangladesh. *BMC Res Notes* 8 (1): 1-9. DOI: 10.1186/s13104-015-1618-6.
- Sofowra A. 1993. *Medicinal Plants and Traditional Medicine in Africa*. Spectrum Books Ltd., Ibadan, Nigeria.
- Tariq AL, Reyaz AL. 2013. Significances and importance of phytochemical present in *Terminalia chebula*. *Intl J Drug Dev Res* 5 (3): 256-262.
- Trease GE, Evans WC. 1989. *Pharmacognosy*, 11th edn., Bailliere Tindall, London.
- Udochukwu U, Omeje FI, Uloma IS, Oseiwe FD. 2015. Phytochemical analysis of *Vernonia amygdalina* and *Ocimum gratissimum* extracts and their antibacterial activity on some drug resistant bacteria. *Am J Res Commun* 3 (5): 225-235.
- Usunobun U, Okolie NP. 2015. Phytochemical, trace and mineral composition of *Vernonia amygdalina* leaves. *Intl J Biol Pharmac Res* 6 (5): 393-399.
- Usunomena U, Ngozi OP. 2016. Phytochemical analysis and proximate composition of *Vernonia amygdalina*. *Intl J Sci World* 4: 11-14. DOI: 10.14419/ijsw.v4i1.5845.
- Wamuyu KR, Machocho AK, Wafula AW. 2020. Antimicrobial and phytochemical screening of *Lannea schweinfurthii* (Engl.) Engl. *Biotechnologi* 17: 1-13. DOI: 10.13057/biofar/c170101.
- Xu R, Zhao W, Xu J, Shao B, Qin G. 1996. Studies on bioactive saponins from Chinese medicinal plants. *Adv Exp Med Biol* 404: 371-82. DOI: 10.1007/978-1-4899-1367-8\_30.

## Effect of palm sap (*Arenga pinnata*) on blood glucose levels of mice (*Mus musculus*) alloxan-induced diabetic

SINDI MARYAM LIPUTO<sup>✉</sup>, DJUNA LAMONDO, MARGARETHA SOLANG

Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Gorontalo. Jl. Prof. BJ. Habibie, Tilongkabila, Bone Bolango 96119, Gorontalo, Indonesia. Tel./fax.: +62-435-821752, ✉email: sindy.liputo@gmail.com.

Manuscript received: 3 January 2022. Revision accepted: 9 May 2022.

**Abstract.** Liputo SM, Lamondo D, Solang M. 2021. Effect of palm sap (*Arenga pinnata*) on blood glucose levels of mice (*Mus musculus*) alloxan-induced diabetic. *Asian J Nat Prod Biochem* 20: 11-15. This study aims to determine the effect of palm sap (*Arenga pinnata* (Wurmb) Merr.) in reducing blood glucose levels in alloxan-induced diabetic mice (*Mus musculus* Linnaeus, 1758). Twenty-five mice aged 2-3 months were grouped into a negative control group, a positive control group, and a palm sap group (0.2 mL, 0.4 mL, and 0.6 mL). The result showed that the palm sap with a dose of 0.4 mL was the most significant in reducing blood glucose levels. This study concluded that the administration of palm sap affected decreasing blood glucose levels ( $p < 0.05$ ).

**Keywords:** Alloxan, blood glucose, mice, palm sap

### INTRODUCTION

Diabetes is one of the fastest-growing health emergencies compared to other diseases. Diabetes mellitus is a condition in metabolism defect that features hyperglycemia symptoms, which require constant medical care. In addition, DM can cause many acute and seriously chronic complications, such as diabetic ketoacidosis, nonketotic hyperosmolar coma, cardiovascular disease, chronic kidney failure, foot ulcers, and eye damage. Based on data from the International Diabetes Federation (IDF), in 2019, people with diabetes reached 465 million adults worldwide, and it is predicted that in 2045 700 million adults will suffer from this disease or an increase of 51% in 2045 (IDF 2019).

As a developing country, Indonesia has experienced increased diabetes mellitus cases. Diabetes affected 7.7 million adults (20-79 years old) in 2013 and 10.7 million adults in 2019. (IDF 2019). According to National Health Survey (Riskesdas) (2018), the number of DM sufferers aged 15 years has increased by 2% compared to 2013. Gorontalo is one of the provinces in the country with the highest number of people with diabetes. Riskesdas (2018) states that the prevalence of diabetes mellitus in Gorontalo has increased to 2.4 percent from only 1.5% in 2013.

Synthetic chemicals and traditional medicines are effective treatments for DM (Diabetes Mellitus) patients. Glibenclamide is one of the synthetic drug classes that are most often used to treat DM because it is widely marketed and the price is affordable. However, glibenclamide has side effects, such as nausea, constipation, dizziness, tremors, and hypoglycemia (Putra et al., 2017). Therefore, as stated by WHO (2004), traditional medicine is used as self-medication or alternative treatment, particularly for chronic diseases such as diabetes and endocrine disorders. Additionally, although WHO recommends traditional

medicine, scientific research is still needed to determine the truth of its efficacy, for example, palm sap (*Arenga pinnata* (Wurmb) Merr.).

Aren is one type of palm plant that is almost spread throughout Indonesia. Almost all parts of this plant can be utilized and used for various needs, both from parts of plant organs (roots, stems, leaves, fruit, etc.), as well as their production (sap, starch/flour) (Lempang 2012; Gunawan et al. 2017). According to Kandowangko et al. (2011), palm sap (*A. pinnata*) has long been utilized as an antidiabetic medicine by the people of Gorontalo, Indonesia. This plant contains water, minerals, carbohydrates, fats, proteins (Ismail et al. 2020), vitamin C (Choong et al. 2016), phenolic, and amino acids (Kurniawan et al. 2018). Palm sap (*A. pinnata*) has several pharmacological, including a diuretic, anti-tuberculosis, and anti-fatigue. In addition, vitamins can help prevent insulin resistance due to Reactive Oxygen Species (ROS) that damage pancreatic cells (Fitriani et al., 2018). Furthermore, given their antioxidant and antihyperglycemic properties, phenolics aid in the reduction of blood glucose in people with diabetes by preventing excessive oxidation and maintaining insulin content in pancreatic cells, binding free radicals and removing them from the body through the excretory system (Wisudanti 2016). Thus, the content of palm sap (*A. pinnata*) has great potential in treating DM.

Palm sap was applied to experimental animals in the form of male mice (*Mus musculus* Linnaeus, 1758). The mice used were normal mice induced with diabetogenic agents. The use of mice as experimental animals is because mice have a high sensitivity compared to other experimental animals to blood glucose tests; male mice are not influenced by hormonal factors such as female mice. In addition, mice have genes similar to humans (Rudiawan 2016). Therefore, before being given palm sap, the mice were first induced with a diabetogenic agent.



Alloxan, a well-known diabetogenic agent, is widely used to induce type 2 diabetes in animals. In addition, alloxan-induced diabetes mellitus serves as a pathological bio model for testing a substance with supposed antioxidant activities. Alloxan can increase blood glucose in mice within 2 x 24 hours without causing death (Rudiawan 2016).

## MATERIALS AND METHODS

### Research design

This study employed a post-test control group design, i.e., the treatment effect, by comparing the treatment group with the control group after being given the action. The experimental design used was a randomized design (CRD) with five treatment groups consisting of two control groups and three groups of palm sap (0.2 mL, 0.4 mL, 0.6 mL).

### Sampling and preparation

Palm sap is obtained from the community in Dulamayo, Bone Bolango District, Gorontalo, Indonesia. Male mice (*M. musculus*) aged 2-3 months were involved as test animals. Alloxan was chosen as the diabetogenic agent since it is the most extensively used chemical for producing diabetes in test animals. Glibenclamide as standard hypoglycemic medication and aquades is utilized as drinking water for mice since it is mineral-free, ensuring that the fall in blood glucose levels is not due to minerals in the water.

### Group of test animals

Mice were divided into five treatment groups; those are K- = negative control group only given aquadest and feed, K+ = positive control group given glibenclamide 0.013 mg/20 g mice BW, K+ = positive control group given glibenclamide 0.013 mg/20 g mice BW, K+ = positive control group given glibenclamide 0.013 mg/20 g mice BW, K+ = positive control group given glibenclamide 0.013 mg/20 g mice BW, K P1 = Treatment 1, palm sap was given at a dose of 0.2 mL/20 g BW mice, P2 = Treatment 2, palm sap was given at a dose of 0.4 mL/20 g BW mice, and P3 = Treatment 3, given palm at a dose of 0.6 mL/20 g BW rats.

### Research procedure

#### *Test animals preparation (acclimatization)*

Mice were acclimatized for seven days to adapt to the new environment and were given water and food ad libitum.

#### *Alloxan treatment to mice*

Alloxan was induced subcutaneously at 100 mg/kg BW dosage after acclimation. Fasting blood glucose (FBG) levels were measured on the third day following alloxan induction. Before measurement, mice fasted for 10 hours. This study employed mice with FBG >126 mg/dL.

### *Test materials provision*

On the 4th day, the group gave glibenclamide and palm sap three days in a row, and the dose was determined after alloxan was induced. Next, GCU measured fasting blood glucose levels on day 6, and the mice fasted for 10 hours before being measured.

### Data analysis

The research data were analyzed using SPSS (Statistical Products and Services Solutions) version 25. Normality was assessed using the Shapiro-Wilk method ( $p > 0.05$ ), and homogeneity was determined using the Levene method ( $p > 0.05$ ) and proceeded with the one-way ANOVA test. If there is a significant difference, it would be continued with Duncan's test to see the difference between the treatment groups.

## RESULTS AND DISCUSSION

### Results


A total of 25 mice (*M. musculus*) were given three different doses of palm sap (0.2 mL, 0.4 mL, and 0.6 mL) and glibenclamide and distilled water as positive and negative controls, respectively. A total of 3 blood draws were performed to measure blood glucose levels (mg/dL). A post hoc test revealed no significant difference between groups ( $p < 0.05$ ) before alloxan induction towards the mice. That is supported by the fact that all groups experienced the same condition. The measurement results of FBG were also within the limits of normal FBG (<126 mg/dL), which shows no hyperglycemic activity in the mice before administering any substance. After the third day of alloxan administration, all groups' FBG showed a hyperglycemic condition with no significant difference. FBG increased significantly ( $p < 0.05$ ) after alloxan administration to mice compared to before administration. After giving palm sap, hyperglycemia was shown in all groups following the decreasing FBG of mice. GDP levels were lower in the palm sap (*A. pinnata*) group (P1, P2, and P3) than in the control group (K- and K+). The data of glucose measurement results can be seen in Table 1 and Figure 1.

Based on the results, the percentage of the palm sap group with a dose of 0.4 mL was more influential in lowering fasting blood glucose in mice after alloxan was induced. An ANOVA statistical test was used to measure if there was a significant effect on each group, and normality and homogeneity checks were performed before the implementation. Based on the normality test results using the Shapiro-Wilk test, the p-value is 0.05, which means the research data is normally distributed. Furthermore, based on the homogeneity test using the Levene test, a p-value of 0.05 was obtained, which means that the fasting blood glucose of mice was homogeneous. Moreover, bearing this in mind, the requirements in the ANOVA test are met.

**Table 1.** Average fasting blood glucose (FBG) levels before and after treatment

Group	Before administration of alloxan $\pm$ SD	After administration of alloxan $\pm$ SD	After administration of palm sap $\pm$ SD	Decrease in FBG after giving palm sap (%)
K- (Without Treatment)	86.20 $\pm$ 3.52 <sup>1.a</sup>	139.80 $\pm$ 3.59 <sup>2.a</sup>	112.80 $\pm$ 3.38 <sup>3.a</sup>	19%
K+ (Glibenclamide)	85.20 $\pm$ 4.64 <sup>1.a</sup>	139.60 $\pm$ 5.05 <sup>2.a</sup>	101.00 $\pm$ 2.07 <sup>3.a</sup>	28%
P1 (Palm Sap 0.2 mL)	81.40 $\pm$ 4.92 <sup>1.a</sup>	133.40 $\pm$ 3.76 <sup>2.a</sup>	84. $\pm$ 5.07 <sup>1.b</sup>	37%
P2 (Palm Sap 0.4 mL)	82.20 $\pm$ 5.28 <sup>1.a</sup>	141.80 $\pm$ 5.81 <sup>2.a</sup>	64.00 $\pm$ 5.32 <sup>3.c</sup>	55%
P3 (Palm Sap 0.6 mL)	81.80 $\pm$ 2.17 <sup>1.a</sup>	134.40 $\pm$ 4.79 <sup>2.a</sup>	78.40 $\pm$ 3.26 <sup>1.b</sup>	42%

Note: Group K- (negative control): alloxan + aquadest, K+ (positive control): alloxan + glibenclamide, P1 (treatment 1): alloxan + palm sap 0.2 mL, P2 (treatment 2): alloxan + palm sap 0.4 mL, P3 (treatment 3): alloxan + palm sap 0.6 mL. The different superscript letters within the same column were significantly different at  $P < 0.05$ . The different superscript numbers within the same row were significantly different at  $P < 0.05$

**Figure 1.** Average fasting blood glucose (FBG) levels in mice before and after treatment

Based on the One Way ANOVA statistical analysis results, there was an effect of giving palm sap on reducing fasting blood glucose in mice with a significant value = 0.00 = 0.05. To see the difference in each treatment entails Duncan's test. Duncan's analysis showed that the K- group was not significantly different from K+ but significantly different from the P1, P2, and P3 groups. The P1 group was significantly different from the K-, K+, and P2 groups but not significantly different from the P3 group. The P2 group differed significantly from the K-, K+, and P1 groups (Table 1).

The normality test utilizing the Shapiro-Wilk test yielded a p-value of 0.05, indicating that the research data is normally distributed. Furthermore, the p-value of 0.05 was achieved based on the Levene test's homogeneity test, indicating that the fasting blood glucose of mice was homogeneous. By this, the ANOVA test's conditions are met. According to the results of a One Way ANOVA statistical analysis, there was an effect of feeding palm sap on reducing fasting blood glucose in mice, with a value of sig = 0.00 = 0.05. Then, I continued Duncan's test to see

how each treatment differed. Duncan's study revealed that the K- group did not differ significantly from the K+ group but considerably from the P1, P2, and P3 groups. The P1 group differed considerably from the K-, K+, and P2 groups but not from the P3 group. The P2 group differed significantly from the K-, K+, and P1 groups (Table 1).

### Discussion

In the fasting blood glucose (GDP) measurement before alloxan was given, all mice were in normal condition (70-100 mg/dL). However, after alloxan was induced, the levels of FBG in mice experienced hyperglycemia (>126 mg/dL). This result proves that alloxan can produce experimental diabetic conditions in test animals. Alloxan is a widely used chemical to induce diabetes in test animals.

Toxicity produced by alloxan occurs due to the formation of free radicals (superoxide radicals and hydroxyl radicals). Free radical activity causes an increase in cytosolic calcium concentration resulting in pancreatic-cell necrosis. A decrease follows this damage in insulin secretion, which results in the reaction of glycogenesis and

reduced glucose transport in cells. In addition, glycogenolysis becomes uncontrolled, resulting in increased blood glucose in the body (Yusni et al., 2017). Meanwhile, Sari et al. (2020) state that alloxan is a toxic glucose analog that selectively destroys insulin-producing pancreatic cells.

In this study, the K- group was the alloxan induction group but was not given any treatment. Before being induced with alloxan, the K- group had a fasting blood glucose of  $86.20 \pm 3.52$  mg/dL (normal); after alloxan was induced, fasting blood glucose increased to  $139.80 \pm 3.59$  mg/dL (hyperglycemia), then there was a decrease in fasting blood glucose by 19%. This decrease is caused by the body's defense efforts to ward off free radicals. In addition, mice's body has endogenous antioxidants such as catalase, superoxide dismutase (SOD), glutathione peroxidase, and glutathione S-transferase (Pratama and Busman 2020).

The K+ group in this study was the group that was given glibenclamide. Before being induced with alloxan, the K+ group had a fasting blood glucose of  $85.2 \pm 4.64$  mg/dL (normal). After alloxan was induced, fasting blood glucose increased to  $139.60 \pm 5.05$  mg/dL (hyperglycemia). Yet, after giving glibenclamide, fasting blood glucose decreased by 28%.

Glibenclamide was administered to the K+ group in this study. Before being induced with alloxan, the K+ group had a fasting blood glucose of  $85.24.64$  mg/dL (normal), fasting blood glucose climbed to  $139.605.05$ mg/dL (hyperglycemia), and fasting blood glucose was reduced by 28% after being given glibenclamide.

Based on the results of the One Way ANOVA test, the most significant group in reducing fasting blood glucose in mice was the palm sap group with a dose of 0.4 mL (P2) and 0.6 mL (P3), but based on the calculation of the percentage of the palm sap group with a dose of 0.4 mL (P2) which most influential with a decrease of 55%. This decrease was higher than the glibenclamide (K+) group of 28%, 0.2 mL of palm juice (P2) 37%, and 0.6 mL of palm juice (P3) 42%. This study was conducted by Swastini et al. (2018), which found that the administration of palm sap reduces fasting blood glucose in male rats induced by alloxan.

The decrease in fasting blood glucose in the palm sap group is thought to be influenced by the compounds contained in palm sap which can prevent the oxidation of pancreatic -cells caused by alloxan induction so that the damage can be controlled. The compounds contained in palm sap are minerals, protein (Ismail et al. 2020), vitamin C (Lempang and Mangopang 2012), triterpenoids, flavonoids, saponins, and alkaloids (Putri et al. 2020).

Protein aids in the regulation of high fasting blood glucose in the body. Based on Gannon et al. (2003), protein can increase insulin concentration. According to Prastari et al. (2017), protein plays a role in inhibiting the work of the -glucosidase enzyme in breaking down carbohydrates into glucose, allowing fasting blood glucose to be managed.

Phenolics play a role in lowering fasting blood glucose by damaging the lipid oxidation structure by donating

hydrogen atoms (Bahman et al. 2019). In addition, phenolic inhibits enzymes involved in carbohydrate metabolism processes, such as -amylase and -glucosidase. Inhibition of this enzyme slows the breakdown of disaccharides into simple glucose so that the body's absorbed glucose is reduced (Oboh and Ademosun 2011). The phenolic groups in palm sap are triterpenoids, flavonoids, saponins, and alkaloids (Putri et al., 2020).

Triterpenoids work as antidiabetics by causing an increase in AMP-activated protein kinase (AMPK) in muscle cells, which enhances glucose consumption in muscle cells and lowers blood glucose. Triterpenoids also help reduce pancreatic-cell damage by delaying the synthesis of TNF- (Tumor Necrosis Factor-alpha) produced by ROS activity (Putra 2013). In insulin-resistant cells, triterpenoids accelerate glucose metabolism and can lower fasting blood glucose in vivo (Hu et al., 2014).

Flavonoids are antioxidants that can inhibit pancreatic cell damage by neutralizing free radicals (Parwata et al., 2018). In addition, flavonoids stop the peroxidation process by inhibiting enzymes that produce superoxide anions and decreasing alkoxy and peroxy radicals (Yuslianti 2017). Furthermore, flavonoids act directly on pancreatic -cells, activating the cAMP signal cascade to boost insulin production, triggered by glucose (Hikmah et al. 2016), preventing glucose absorption in the gut (Parwata et al. 2018).

Alkaloids have therapeutic effects and are antidiabetic since these compounds have the potential to interact with various proteins involved in glucose homeostasis (Rasouli et al., 2020). Alkaloids act as antidiabetics, both in extracts and isolated molecules. It inhibited the glucosidase enzyme, deactivated DPP-IV (Dipeptidyl Peptidase-4), improved insulin sensitivity, and reduced oxidative stress (Ajebli et al., 2021)

Vitamin C helps repair damaged cells, insulin resistance, and inflexible blood vessels associated with diabetes (Aprila et al. 2015). Vitamin C is a group of micronutrients that plays a role as an antioxidant in human plasma and functions as an electron donor or electron-reducing agent, which allows it to serve as an antioxidant. According to Maha (2013), Vitamin C can lower fasting blood glucose, providing primary and secondary protection against oxidative damage to lipids and lipoproteins and improving the negative effects of DM.

Aprila et al. (2015) stated that magnesium simplifies glucose transport into cells, cofactors various enzymes involved in the glucose oxidation process, and helps to increase insulin sensitivity and avoids insulin resistance (Larsson and Wolk 2007). In addition, intracellular magnesium deficiency can reduce the action of tyrosine kinase on insulin receptors (Mulatsih 2020).

In conclusion, the administration of palm sap (*A. pinnata*) can lower fasting blood glucose in mice (*M. musculus*) with alloxan-induced diabetes. Palm sap (*A. pinnata*) 0.4 mL was the most significant in lowering the mice's fasting blood glucose.

## REFERENCES

- Ajebli M, Khan H, Eddouks M. 2021. Natural alkaloids and diabetes mellitus: A review. *Endocr Metab Immun Disord Drug Targets* 21 (1): 111-130. DOI: 10.2174/1871530320666200821124817.
- Aprila IF, Salim MN, Daud R, Armansyah T, Asmilina N, Jamin F. 2015. Pengaruh pemberian kacang panjang (*Vigna unguiculata*) terhadap kadar glukosa darah mencit (*Mus musculus*) yang diinduksi aloksan. *J Medika Veterenia* 9(2): 97-99. DOI: 10.21157/j.med.vet.v9i2.3983. [Indonesian]
- Bahman DS, Yuliet, Ihwan. 2019. Efek akar *Garcinia rostrata* Hassk.ex Hook.f terhadap penurunan kadar glukosa darah puasa mencit jantan (*Mus musculus*) yang diinduksi aloksan. *Biocelebes* 13 (1): 21-29. [Indonesian]
- Choong CC, Anzian A, Sapawi CW, Hussin M. 2016. Characterization of Sugar from *Arenga pinnata* and *Saccharum officinarum* Sugars. *Intl Food Res J* 23 (4): 1642-1652.
- Fitriani LI, Murbawani E, Nissa C. 2018. Hubungan asupan vitamin C, vitamin E dan B-karotendengan kadar gula darah puasa pada wanita usia 35-50 tahun. *J Nutr Collage* 7 (2): 84-91. DOI: 10.14710/jnc.v7i2.20827. [Indonesian]
- Gannon MC, Nuttal FQ, Saeed S, Jordan K, Hoover H. 2003. An increase in dietary protein improves the blood glucose response in persons with type 2 diabetes. *Am J Clin Nutr* 78: 734-741. DOI: 10.1093/ajcn/78.4.734.
- Gunawan R, Ramadhan UG, Iskandar J, Partasmita R. 2017. Local knowledge of utilization and management of sugar palm (*Arenga pinnata*) among Cipanggulaan People of Karyamukti, Cianjur (West Java, Indonesia). *Biodiversitas* 19: 93-105. DOI: 10.13057/biodiv/d190115.
- Hikmah N, Yuliet, Khaerati K. 2016. Pengaruh pemberian ekstrak daun salam (*Syzygium polyanthum* Wight.) terhadap glibenklamid dalam menurunkan kadar glukosa darah mencit (*Mus musculus*) yang diinduksi aloksan. *Galenika J Pharm* 2 (1): 24-30. DOI: 10.22487/j24428744.2016.v2.i1.5300. [Indonesian]
- Hu X, Wang S, Xu J, Wang D-B, Chen Y, Yang G-Z. 2014. Triterpenoid saponins from *Stauntonia chinensis* Ameliorate insulin resistance via the AMP-Activated Protein Kinase and IR/IRS-1/PI3K/Akt pathways in insulin-resistant HepG2 cells. *Intl J Mol Sci* 15: 10446-10458. DOI: 10.3390/ijms150610446.
- International Diabetes Federation (IDF). 2019. IDF Diabetes Atlas. Diakses 18 januari 2020 [www.diabetesatlas.org](http://www.diabetesatlas.org) [Indonesian]
- Ismail, Yurnia NN, Solang M, Uno WD. 2020. Komposisi proksimat dan indeks glikemik nira aren. *Biospecies* 13 (2): 1-9. DOI: 10.22437/biospecies.v13i2.8761. [Indonesian]
- Kandowangko, Novri Y, Solang M, Ahmad J. 2011. Kajian Etnobotani Tanaman Obat oleh Masyarakat Kabupaten Bonebolango Provinsi Gorontalo. Laporan Penelitian Etnobotani Tanaman Obat Jurusan Biologi Universitas Negeri Gorontalo, Gorontalo. [Indonesian]
- Kurniawan T, Jayanudin, Kustiningsih I, Firdaus MA. 2018. Palm sap sources, characteristic, and utilization in Indonesia. *J Food Nutr Res* 6 (9): 590-596. DOI: 10.12691/jfnr-6-9-8.
- Larsson SC, Wolk A. 2007. Magnesium intake and risk of type 2 diabetes: A meta-analysis. *J Intern Med* 262: 208-214 DOI: 10.1111/j.1365-2796.2007.01840.x.
- Lempang M, Mangopang AD. 2012. Efektivitas nira aren sebagai bahan pengembang adonan roti. *J Penelitian Kehutanan Wallacea* 1 (1): 26-35. DOI: 10.18330/jwallacea.2012.vol1iss1pp26-35. [Indonesian]
- Lempang M. 2012. Pohon aren dan manfaat produksinya. *Info Teknis Eboni* 9 (1): 37-54. [Indonesian]
- Maha MH. 2013. Potential of vitamin C and/or *Lupinus termis* to modulate blood glucose level and oxidative stress status of alloxan-induced diabetic rats. *Sci J Biol Sci* 2 (11): 41-46.
- Mulatsih ES. 2020. Hubungan Asupan Magnesium dengan Kadar Glukosa Darah pada Penderita Diabetes Mellitus Tipe II Di Puskesmas Tawang Sari Sukoharjo. [Hon. Thesis]. Fakultas Ilmu Kesehatan, Universitas Muhammadiyah Surakarta. [Indonesian]
- Oboh G, Ademosun AO. 2011. Shaddock peels (*Citrus maxima*) phenolic extracts inhibit  $\alpha$ -amylase,  $\alpha$ -glucosidase and angiotensin I-converting Enzym Activities: Anutraceutical approach to diabetes management. *Diabetes Metab Syndr* 5: 148-152. DOI: 10.1016/j.dsx.2012.02.008.
- Parwata A, Laksmiwati, Dina MN, Yasa S. 2018. The contents of phenol and flavonoid compounds in water extract of *Gyrinops versteegii* leaves have potential as natural antioxidants and hypoglycemic in hyperglycemic Wistar rats. *Biochem Pharmacol* 11 (3): 1543-1552. DOI: 10.13005/bpj/1521.
- Prastari C, Yasni S, Nurilmala M. 2017. Karakteristik protein ikan gabus yang berpotensi sebagai antihiperlipidemik. *JPHPI* 20 (2): 413-423. DOI: 10.17844/jphpi.v20i2.18109. [Indonesian]
- Pratama NA, Busman H. 2020. Potensi antioksidan kedelai (*Glycine Max L*) terhadap penangkapan radikal bebas. *J Ilmiah Kesehatan Sandi Husada* 11 (1): 497-504. DOI: 10.35816/jiskh.v11i1.333. [Indonesian]
- Putra AP. 2013. Pengaruh Pemberian Ekstra Etanol Buncis (*Phaseolus vulgaris* L.) terhadap Penurunan Kadar Glukosa Darah Menicit yang Diinduksi Aloksan. [Hon. Thesis]. Fakultas Kedokteran, Universitas Jember, Jember. [Indonesian]
- Putra, Achmad A, Rachma H. 2017. Kejadian efek samping potensial terapi obat anti diabetes pasien diabetes melitus berdasarkan Algoritma Naranjo. *Pharm J Indonesi* 2 (2): 45-50. DOI: 10.21776/ub.pji.2017.002.02.3. [Indonesian]
- Putri IDAEW, Ratnayanti IGAD, Sugiritma IW, Arijana IGKN. 2020. Analisis fitokimia nira aren dan tuak aren (*Arenga pinnata* (Wurmbr) Merr.). *J Medika Udayana* 10 (6): 18-22. [Indonesian]
- Rasouli H, Yarani R, Paciot F, Popovic-Djordjevic J. 2020. Antidiabetic potential of plant alkaloids: Revisiting current findings and future perspectives. *Pharmacol Res* 155: 104723. DOI: 10.1016/j.phrs.2020.104723.
- Riskesdas. 2018. Laporan Nasional Riskesdas 2018. Balitbangkes. Accessed January 18, 2020 [www.labdata.litbang.kemkes.go.id](http://www.labdata.litbang.kemkes.go.id)
- Rudiawan A. 2016. Pengaruh Aktivitas Fisik Maksimal terhadap Peningkatan Jumlah Leukosit Pada Tikus Putih (*Rattus Norvegicus* Wistar). [Diploma Thesis]. Universitas Negeri Malang, Malang. [Indonesian]
- Sari ADP, Kusuma H, Ratri DMN, Wibisono C, Suprapti B. 2020. The effect of premixed insulin to blood glucose concentration in patients with type 2 diabetes mellitus. *J Basic Clin Physiol Pharmacol* 2020: 20190342. DOI: 10.1515/jbcpp-2019-0342.
- Wisudanti DD. 2016. Aplikasi terapeutik geraniin dari ekstrak kulit rambutan (*Nephelium lappaceum*) sebagai antihiperlipidemik melalui aktivitasnya sebagai antioksidan pada diabetes melitus tipe 2. *Nurseline J* 1 (1): 120-138. [Indonesian]
- World Health Organization (WHO). 2004. Guidelines for the Regulation of Herbal Medicines in the South-East Asia Region. accessed on 17 November 2020 <https://apps.who.int/iris/handle/10665/205175>
- Yuslianti ER. 2017. Pengantar Radikal Bebas dan Antioksidan. Deepublish, Yogyakarta. [Indonesian]
- Yusni, Akbar IB, Rezanita, Fahlevi R. 2017. Penurunan kadar gula darah akibat pemberian ekstrak manggis (*Gracinia mangostana*) dan tomat (*Lycopersicon esculentum* Mill) pada tikus diabetes. *Glob Med Health Commun* 5 (1): 57-63. DOI: /10.29313/gmhc.v5i1.2097.

## Short Communication: Extraction and characterization of pectin from ripe and unripe mango (*Mangifera indica*) peel

CHRISTOPHER O. SHAIBU\*, JULIUS DINSHIYA, VIVIAN EBERE SHAIBU

Department of Biochemistry, Federal University Wukari. PMB 1020 Katsina-Ala Road, Wukari, Taraba State, Nigeria. Tel.: +2348030000000,

\*email: chrishaibu@fuwukari.edu.ng

Manuscript received: 6 April 2022. Revision accepted: 12 June 2022.

**Abstract.** Shaibu CO, Dinshiya J, Shaibu VE. 2022. Short Communication: Extraction and characterization of pectin from ripe and unripe mango (*Mangifera indica*) peel. *Asian J Nat Prod Biochem* 20: 16-20. Pectin is a heteropolysaccharide present in the cell walls of different plants at different concentrations with widespread applications. This work aimed at extracting and characterizing pectin from ripe and unripe mango peel to investigate the effect of ripeness on the yield and quality of mango pectin. Pectin was extracted at varying temperatures, time, and pH using 0.1 N HCl as the extraction solvent to obtain the optimum extraction condition. The maximum yield of pectin was 22.67% for ripe mango peel and 21.90% for unripe mango peel. The optimum extraction conditions were 90°C, 60 mins, and pH 1.5. The pectin extracted using the optimum extraction conditions was then characterized. The moisture content, ash content, methoxyl content, equivalent weight, anhydrouronic acid, and degree of esterification of the ripe mango peel pectin were found to be 8.76±0.08%, 10.12±0.47%, 9.17±0.27%, 883.07±13.85 g, 72.45±0.59 and 72.52±0.09%, respectively. In contrast, those of unripe mango peel pectin was found to be 8.13±0.13%, 9.12±0.34%, 8.83±0.19%, 823.38±14.07 g, and 71.56±0.34%, 70.34±0.38%, respectively. This study showed that ripe mango peel might be more suitable for use as a pectin source. However, pectin extracted from ripe and unripe mango peel could be considered an alternative source of pectin in food processing, pharmaceutical industries, and various places of pectin application.

**Keywords:** Characterization, extraction, mango, pectin, yield

### INTRODUCTION

According to research by Singh et al. (2021), over 2 billion tonnes of agro-waste are generated annually worldwide. This volume of waste is capable of causing environmental and health hazards. Therefore, their reuse has been strongly advocated to avert the undesirable effects of their accumulation on the environment and maximize benefits from their utilization (Harshwardhan and Upadhyay 2017).

Mango (*Mangifera indica* L.) belongs to the family Anacardiaceae with over 1,365 varieties worldwide (Mubarik et al. 2020). Mangoes have been reported to contain high vitamin C, fiber, and pectin (Mubarik et al., 2020). In 2017, Mango production was reported to be about 47 million tonnes per annum (Altendorf 2017). In addition, 40-50% of the fruit constitutes agro-waste (Rai et al. 2020) which can serve as a good source for pectin production.

Pectin belongs to the family of heteropolysaccharides primarily found in the primary cell walls of terrestrial plants (Maxwell et al. 2012; Hamed and Mustafa 2018; Serna and Ayala 2020). It is constituted of 4 polymers rich in D-galacturonic acid (GalA) and containing significant amounts of L-rhamnose (Rha), D-arabinose (Ara), and D-galactose (Gal) (Fissore et al. 2012).

Agro-industrial wastes utilized for pectin production include fruit wastes such as sweet melon (Rahmani et al. 2020), kiwifruit pomace (Yuliarti et al. 2015), papaya peels

(Koubala et al. 2014), lemon, grapefruit, orange, avocado, (Bamba et al. 2020), sugar beet (Maxwell et al. 2016).

Pectin is highly valued as a functional food ingredient and has wide pharmaceutical/medicinal applications, including drug delivery, gene delivery, wound healing, and tissue engineering (Maran et al., 2013; Wicker et al., 2014).

Extraction of pectin usually involves hydrolysis of pectin macromolecules under controlled temperature, pH, and time. Temperature, pH, extraction solvent, and extraction time have significantly affected pectin yield (Kliemann et al., 2009; Sharma et al., 2013). Acid extraction seems to be the most widely used method, although other methods like the use of enzymes, microwave, subcritical water, and high pressure have been employed in the extraction of pectin (Zoghi et al., 2021)

This study aimed at extracting and characterizing pectin from ripe and unripe mango peel to investigate the effect of ripeness on pectin yield and quality from mango.

### MATERIALS AND METHODS

#### Materials

All the chemicals used were of analytical grade and included Sodium chloride (NaCl), distilled water, ethanol (C<sub>2</sub>H<sub>5</sub>OH), Hydrochloric Acid (HCl), phenol red, and sodium hydroxide (NaOH).

## Equipment

Mortar and pestle, Beakers, Conical flasks, Spatula, Filter paper (Whatman No. 1), Masking tape, Separator funnel, Digital analytical weighing balance, Spectrophotometer (Jenway: 6305), Refrigerator, Thermostatic Water Cabinet (Model: HH - W420), Micropipette, Test tubes and Test Tube Racks, Centrifuge (Model: 80-2B), and Sample Bottles.

## Sample collection

Both ripe and unripe Mango fruit were collected from Federal University Wukari farmland, Taraba state, Nigeria, located on 7°50'34.0" N 9°46'19.1" E.

## Methods

### Sample preparation

Ripe and unripe mango fruits (*M. indica*) were peeled, chopped, and blanched in a water bath at 80°C for 5 minutes. The blanched sample was then oven-dried at 60°C for 72 hours and ground into fine powder. It was then sieved through a 60mm to 80mm mesh sieve and stored in a separate air-tight container until it was required for further analysis.

**Note:** The degree of ripeness of the mango used in this study was 4 for unripe and 8 for ripe, using a scale of 1 to 10, with 1 as the most unripe and 10 as the ripest.

### Pectin extraction

Pectin extraction was performed using 0.1 N HCl solvent at different extraction temperatures (70, 80, and 90°C), extraction times (30, 60 and 90 mins), and pH (1.5, 2.5, and 3.0). 10 g of the powdered ripe mango peel was macerated in 250 mL extraction solvent, and the pH was adjusted with 0.1 N HCl and NaOH. Afterward, the mixture was heated in a water bath for the specified time. The mixture was filtered through a cheesecloth and pressed to recover the extract; pectin was then precipitated from the mixture using 98% ethanol in a ratio of 1:2 (1 part extract and 2 parts ethanol) and kept at room temperature overnight. The precipitated pectin was then filtered through Whatman No.1 (filter paper) and washed with 70% ethanol (v/v) and 80% ethanol (v/v) to remove the soluble impurities. The extracted pectin was then dried at 60°C for 24 hours in an oven. The same procedure was repeated for unripe mango peel.

## Determination of % yield

Pectin yield was calculated using the formula:

$$\frac{\text{Weight of pectin}}{\text{Weight of sample used}} \times \frac{100}{1}$$

### Pectin characterization

**Ash content:** Ash content of pectin was determined by Ranganna's method (Ranganna 1986). A 1.2 g pectin (sample) was weighed, the sample was ignited slowly in a crucible and then heated for 3-4 hours at 600°C, then the crucible was taken into the desiccators and allowed to cool.

Ash content was determined using the equation:

$$\text{Ash\%} = \frac{\text{Weight of ash}}{\text{Weight of sample}} \times 100$$

**Moisture content:** The moisture content of pectin was determined by Ranganna's method (Ranganna 1986). A 1 g pectin sample was weighed and placed into a metal dish. The sample was dried in an oven for 5 hours at 100°C, cooled in a desiccator, and weighed.

The moisture content was determined using the equation:

$$\text{Moisture content (\%)} = \frac{\text{Weight of residue}}{\text{Weight of sample}} \times 100$$

**Equivalent weight:** Equivalent weight was determined by Ranganna's method (Ranganna 1986). A 0.5 g pectin was measured into a 250 mL conical flask, a 5 mL ethanol was added, and 1 g of sodium chloride and 100 mL of distilled water were added to the solution, after which 6 drops of phenol red were added. The mixture was titrated against 0.1 N NaOH. A purple color indicated the titration endpoint. The neutralized solution was stored for the determination of methoxyl content and anhydrouronic acid content.

$$\text{Equivalent Weight} = \frac{\text{Weight of sample} \times 1000}{\text{ml of alkali} \times \text{Normality of alkali}}$$

**Methoxyl content (MeO):** The methoxyl content is an important factor in controlling the setting time of pectins. MeO was determined using Ranganna's method (Ranganna 1986). First, the 25 mL sodium hydroxide (0.25 N) was added to the neutral solution obtained from the determination of equivalent weight. Next, the solution was stirred thoroughly and kept at room temperature for 30 mins. After 30 min, 25 mL of 0.25 N hydrochloric acid was added and titrated against 0.1 N NaOH.

Methoxyl content was calculated by using the following formula:

$$\text{Methoxyl content\%} = \frac{\text{ml of alkali} \times \text{Normality of alkali} \times 3.1}{\text{Weight of sample}}$$

**Total anhydrouronic acid content (AUA):** Total AUA of pectin was obtained mathematically using the following formula:

$$\text{AUA} = \frac{176 \times 0.1z \times 100}{W \times 1000} + \frac{176 \times 0.1y \times 100}{W \times 1000}$$

Where: Molecular unit of AUA (1 unit) = 176 g, z = mL (titre) of NaOH from equivalent weight determination, y = mL (titre) of NaOH from methoxyl content determination and w = weight of sample.

### Degree of esterification (DE)

The DE of pectin was measured based on methoxyl content (MeO) and AUA and calculated using the formula:

$$\text{Degree of esterification (DE) \%} = \frac{176 \times \% \text{ MeO}}{31 \times \% \text{ AUA}} \times 100$$

### Statistical analysis

All data are represented as means  $\pm$  SD. Statistical analyses were performed using the Independent samples t-test on Statistical Package for Social Sciences (SPSS) version 20.

## RESULTS AND DISCUSSION

### Percentage (%) yield of pectin at pH 1.5, 60 minutes, and varying temperature

The percentage yield of pectin at 60 mins, pH 2.0, and different temperatures are shown in Figure 1. Although the percentage yield of pectin increased with an increase in temperature for both mango peels, pectin from the ripe mango was higher than that of the unripe mango peel.

### Percentage (%) yield of pectin at 90°C, 60 minutes, and varying pH

The percentage yield of pectin at 60 mins, temperature 90°C, and different pHs of 1.5, 2.0, and 3.0 are shown in Figure 2. Pectin yield for ripe and unripe mango peel decreased with an increase in pH. The yield of the ripe mango peel was higher than that of the unripe mango peel.

### Percentage (%) yield of pectin at 90°C, pH 1.5, and varying time


Figure 3 shows the percentage yield of pectin at 90°C, pH 1.5, and at different times. There was an increase in

pectin yield for both ripe and unripe mango peel from 30 mins to 60 mins extraction time. However, there was a decrease from 60 mins to 90 mins. The pectin yield of the ripe mango peel was higher when compared to that of the unripe mango peel.


## Discussion

### Pectin yield


This study considered pectin extraction from ripe and unripe mango peel to investigate the effect of ripeness on the yield and quality of pectin produced from mango. Pectin was extracted at various temperatures (70°C, 80°C, and 90°C), pH (1.5, 2.0, and 3.0), and time (30 mins, 60 mins, and 90 mins) to obtain the optimum extraction condition. The optimum extraction conditions shown in this study were pH 1.5, 60 mins extraction time, and 90°C. Pectin extracted using the optimum extraction condition was used in characterization. Studies on the effect of pH, temperature, and time showed that the total yield of pectin for both ripe and unripe mango peel increased with a decrease in pH from pH 3.0 to pH 1.5. That agrees with the work of Hamidon and Zaidel (2017), who reported that the yield of pectin increased with a decrease in the pH of the extractant while working on potato peel residue. Pectin yield also increased with extraction time; however, further increasing the extraction time to 90mins showed a decrease in pectin yield. Thus, the optimum time of the extraction for the maximum yield of mango peel pectin was found to be 60 mins. Similar results were reported on pectin extracted from mandarin orange peels by Koubala et al. (2008). The effect of extraction temperature on the yield is shown in Figures 2 and 3. The yield of pectin increased significantly with the increase in extraction temperature. Similar observations were reported for dried mango peel (Sangheetha et al., 2018).



**Figure 1.** % Yield of pectin at pH 1.5, 60 minutes, and varying temperature



**Figure 2.** % Yield of pectin at 90°C, 60 minutes, and varying pH



**Figure 3.** % Yield of pectin at 90°C, pH 1.5, and varying time

The maximum yield was 22.67% and 21.94% for ripe and unripe mango peel, respectively. Higher pectin yield for ripe mango peel may result from a higher concentration of pectin due to the maturation of the fruit. On the other hand, pectin concentration may decrease with an increase in ripeness due to pectin hydrolysis by pectinase (Prasanna et al., 2007). That is as opposed to the findings of Castillo-Israel et al. (2015), who had higher values for pectin yield from an unripe banana than from a ripe banana peel. However, the yield of pectin from both ripe and unripe mango peel compares well to the 2.93-25.80% pectin yield recorded by Bamba et al. (2020) from orange, lemon, grapefruit, and avocado and is higher than those reported by Yapó and Koffi (2006), for passion fruit pectin (7.5%), and ambarella peel pectin (10-13%).

### Characterization of mango peel pectin

Ripe mango pectin had significantly higher values for yield moisture content, ash content, equivalent weight, and degree of esterification (Table 1). There was no significant difference in methoxyl content and anhydrouronic acid. The moisture content of ripe and unripe mango peel pectin compares well with 10.59 and 11.68 reported for mango peel and watermelon pectin, respectively, by Rury et al. (2017). However, the moisture content of the unripe mango pectin was significantly lower than that of the ripe mango peel.

Ash content measures the total amount of minerals present within a food. The ash content of the pectin isolated in this work was as high as 9-11%. Nazaruddin et al. (2011) reported that the maximum value needed for the quality gel is 10%. That shows that pectin extracted from both mango peels could be a good source for producing quality gel.

There was no significant difference in the methoxyl contents of pectin extracted from ripe and unripe mango peel. The Methoxyl content of commercial pectins generally varies from 8-11% and can form high-sugar gels (>65% sugar) (Castillo-Israel et al. 2015). Therefore, this study's values obtained from ripe and unripe mango peel compare favorably with commercially available pectin.

The extracted pectin's equivalent weight was  $883.07 \pm 13.85$  g and  $823.38 \pm 14.07$  g for ripe and unripe mango peel, respectively. However, this indicates that the ripe mango pectin has a significantly higher gel-forming ability than the unripe. Values obtained in this study were lower than those of apple pomace pectin (833.33-1666.30), Kumar and Chauhan (2010), but higher than those of cocoa husk pectin (510.68-645.19), as reported by Ramli and Asmawati (2011).

Anhydrouronic acid (AUA) is essential to determine the purity and degree of esterification and evaluate physical properties (Ranganna 1986). It indicates the purity of extracted pectin. Pectin with an AUA of 65% or more is considered pure (Nazaruddin et al. 2011). Values for AUA (%) obtained in this study are greater than 65%, indicating that they are pure.

This study's degree of esterification (DE) was within the 60-90% range, generally found in plant tissues (Shaha et al., 2013). Pectins could be classified as rapid-set (DE >72%) and slow-set (DE: 58-65%). That describes the rate of gel formation. Pectin from ripe and unripe mango peels had DE values less than 75% but greater than the range for slow-set pectins; this shows that they may be classified as rapid-set pectin. Values obtained were lower than those reported by Wang et al. (2014), who reported DE values of 76.30 and 83.41% in pectin from the fruit peel of *Citrus maxima* and apple pomace, respectively.

**Table 1.** Pectin characterization

Parameter	Ripe (%)	Unripe (%)	<sup>A</sup> Mean differences	<sup>P</sup> values of mean differences
Yield	22.67	21.90	0.77	0.002 <sup>c</sup>
Moisture content	8.76±0.08	8.13±0.13	0.63	0.021 <sup>c</sup>
Methoxy content	9.17±0.27	8.83±0.19	0.34	0.668 <sup>b</sup>
Ash content	10.12±0.47	9.12±0.34	1.00	0.026 <sup>c</sup>
Equivalence weight	883.07±13.85g	823.38±14.07	59.69	0.003 <sup>c</sup>
Anhydrouronic acid	72.45±0.59	71.56±0.34	0.89	0.148 <sup>b</sup>
Degree of esterification	72.52±0.09	70.34±0.38	2.18	0.007 <sup>c</sup>

Note: Values represent Mean ± standard deviation. <sup>a</sup>= indicates the mean values of parameters for ripe seeds minus the mean values of unripe ones. <sup>b</sup>= Mean difference is not significant <sup>c</sup>=Mean difference is significant at P<0.05



The equivalent weights of the extracted pectin were found to be high, indicating that they have a low partial degradation property. The degree of esterification, equivalent weight, and ash content of ripe mango pectin were significantly higher than that of the unripe mango peel pectin. There was no significant difference in methoxyl and anhydrouronic acid content, thus indicating that the quality of pectin obtainable from ripe mango peel is comparatively better than pectin obtained from unripe mango peel in terms of gel formation and purity. However, both ripe and unripe mango peel pectin is comparable to apple pomace and citrus, thus, indicating the possible use of mango peel as an alternative source in commercial pectin production and for industrial, pharmaceutical, medical, and agricultural applications.

## REFERENCES

- Altendorf S. 2017. Global prospects for major tropical fruits. Short term outlook, challenges and opportunities in a vibrant global market place. *Food Outlook* 2017: 69-81.
- Bamba SS, Gouin EJ, Kouassi AC, Komenan MS, Akre H, Soro YR. 2020. Production of pectin as a relevant tool for by-products management resulting from four tropical edible fruits processing: Extraction yield, physicochemical and functional properties. *Intl J Chem Process Eng Res* 7: 60-73. DOI: 10.18488/journal.65.2020.71.60.73.
- Castillo-Israel KAT, Baguio SF, Diasanta MDB, Lizardo RCM, Dizon EI, Mejico, MIF. 2015. Extraction and characterization of pectin from Saba banana [*Musa 'saba' Musa acuminata* x *Musa balbisiana*] peel wastes" A preliminary study. *Intl Res J* 221: 202-207.
- Fissore EN, Rojas AM, Gerschenson LN. 2012. Rheological performance of pectin enriched products isolated from red beet *Beta vulgaris* L. var. *conditiva* through alkaline and enzymatic treatments. *Food Hydrocoll* 26: 249-260. DOI: 10.1016/j.foodhyd.2011.06.004.
- Hamed AAR, Mustafa SE. 2018. Extraction and assessment of pectin from pumpkin peels. *Biofarmasi J Nat Prod Biochem* 16: 1-7. DOI: 10.13057/biofar/f160101.
- Hamidon NH, Zaidel DNA. 2017. Effect of extraction conditions on pectin yield extracted from potato peels residue using hydrochloric acid. *Chem Eng Trans* 56: 979-984. DOI: 10.3303/CET1756164.
- Harshwardhan K, Upadhyay K. 2017. Effective utilization of agricultural waste: A review. *J Fundamen Renew Energ Appl* 7: 237. DOI: 10.4172/20904541.1000237.
- Kliemann E, Simas KN, Amante ER, Schwinden PE, Reinaldo FT, Marcia MCF, Renata DM. 2009. Optimisation of pectin acid extraction from passion fruit peel (*Passiflora edulis* flavicarpa) using response surface methodology. *Intl J Food Sci Technol* 44: 476. DOI: 10.1111/j.1365-2621.2008.01753.x.
- Koubala BB, Christiaens S, Kansci G, Loey AMV, Hendrickx ME. 2014. Isolation and structural characterizations of papaya peel pectin. *Food Res Intl* 55: 215-221. DOI: 10.1016/j.foodres.2013.11.009.
- Koubala BB, Mbome LI, Kansci G, Mbiap TF, Crepeau MJ, Thibault, F, Ralet MC. 2008. Physicochemical properties of pectins from ambarella peels *Spondias cytherea* obtained using different extraction conditions. *Food Chem* 106: 1202-1207. DOI: 10.1016/j.foodchem.2007.07.065.
- Kumar A, Chauhan GS. 2010. Extraction and characterization of pectin from apple pomace and its evaluation as lipase steapsin inhibitor. *Carbohydr Polym* 82: 454-459. DOI: 10.1016/j.carbpol.2010.05.001.
- Maran JP, Sivakumar V, Thiruganasambandham K, Sridhar R. 2013. Optimization of microwave assisted extraction of pectin from orange peel. *Carbohydr Polym* 97: 703-709. DOI: 10.1016/j.carbpol.2013.05.052.
- Maxwell EG, Belshaw NJ, Waldron KW, Morris VJ. 2012. Pectin-An emerging new bioactive food polysaccharide. *Trends Food Sci Technol* 24: 64-73. DOI: 10.1016/j.tifs.2011.11.002.
- Maxwell EG, Colquhoun IJ, Chau HK, Hotchkiss AT, Waldron KW, Morris VJ, Belshaw NJ. 2016. Modified sugar beet pectin induces apoptosis of colon cancer cells via an interaction with the neutral sugar side-chains. *Carbohydr Polym* 136: 923-929. DOI: 10.1016/j.carbpol.2015.09.063.
- Mubarik F, Noreen S, Farooq F, Siddiq A, Khan M. 2020. A review on pharmaceutical and nutritional benefits of mango *Mangifera indica* Linn: A remedy for cancer, gastrointestinal infections. *Abasyn J Life Sci* 32: 82-92. DOI: 10.34091/AJLS.3.2.8.
- Nazaruddin R, Norazelina SMI, Norziah MH, Zainudin M. 2011. Pectins from dragon fruit *Hylocereus polyrhizus* peel. *J Malay Appl* 401: 19-23.
- Prasanna V, Prabha TN, Tharanathan RN. 2007. Fruit ripening phenomena-an overview. *Crit Rev Food Sci Nutr* 47 (1): 1-19. DOI: 10.1080/10408390600976841.
- Rahmani Z, Khodaiyan F, Kazemi M, Sharifan A. 2020. Optimization of microwave assisted extraction and characterization of pectin from sweet lemon peel. *Intl J Biol Macromol* 147: 1107-1115. DOI: 10.1016/j.jbiomac.2019.10.079.
- Rai AK, Das H, Dash S, Behera S. 2020. Mango seed: A potential source of nutrition from waste. *Indian Farmer* 7: 24-248.
- Ramli N, Asmawati S. 2011. Effect of ammonium oxalate and acetic acid at several extraction time and pH on some physicochemical properties of pectin from cocoa husk *Theobroma cacao*. *Afr J Food Sci* 5: 790-798. DOI: 10.5897/AJFS11.107.
- Ranganna S. 1986. *Handbook of Analysis and Quality Control for Fruit and Vegetable Products*. Tata Mc Graw-Hill Publishing Company, New Delhi.
- Rury IA, Mohamad, Siti S. 2017. Characteristics of pectin isolated from mango (*Mangifera indica*) and watermelon (*Citrullus vulgaris*) peel. *J Appl Food Technol* 4 (2) 31-34. DOI: 10.17728/jaft.42.
- Sangheetha S, Illeperuma DCK, Navaratne AN, Jayasinghe C. 2018. Effect of pH, temperature and time combination on yield and degree of esterification of mango peel pectin: A Box-Behnken design based statistical modelling. *Trop Agric Res* 302: 1-12. DOI: 10.4038/tar.v30i2.8304.
- Serna SO, Ayala SFE. 2020. Chemical composition and biosynthesis of dietary fiber components. In: Welti-Chanes J, Serna-Saldívar S, Campanella O, Tejada-Ortigoza V (eds). *Science and Technology of Fibers in Food Systems*. Food Engineering Series. Springer, Cham. DOI: 10.1007/978-3-030-38654-2\_2.
- Shaha RK, Nayagi Y, Punichelvana A, Afandi A. 2013. Optimized extraction condition and characterization of pectin from Kaffir lime *Citrus hystrix*. *Malaysia: Res J Agric For Sci* 12: 1-11.
- Sharma H, Bhatia, S, Alam MS. 2013. Studies on pectin extraction from kinnow peel and pomace. *J Agric Res* 50: 128.
- Singh R, Das R, Sangwan BT, Khanam R, Pedda GP, Das S, Misra S. 2021. Utilization of agro-waste for sustainable green production: A review. *Environ Sustain* 4: 619-636. DOI: 10.1007/s42398-021-00200-x.
- Wang X, Chen Q, Lü X. 2014. Pectin Extracted from apple pomace and citrus peel by subcritical water. *Food Hydrocoll* 38: 129-137. DOI: 10.1016/j.foodhyd.2013.12.003.
- Wicker L, Kim Y, Kim M, Thirkield B, Lin Z, Jung J. 2014. Pectin as a bioactive polysaccharide-extracting tailored function from less. *Food Hydrocoll* 42: 251-259. DOI: 10.1016/j.foodhyd.2014.01.002.
- Yapo BM, Koffi KL. 2006. Yellow passion fruit rind e a potential source of low-methoxyl pectin. *J Agric Food Chem* 54: 2738-2744. DOI: 10.1021/jf052605q.
- Yuliarti O, Goh KKT, Matia-Merino L, Mawson J, Brennan C. 2015. Extraction and characterisation of pomace pectin from gold kiwifruit *Actinidia chinensis*. *Food Chem* 187: 290-296. DOI: 10.1016/j.foodchem.2015.03.148.
- Zoghi A, Vedadi S, Esfahani ZH, Gavlighi HA, Khosravi D. 2021. A review on pectin extraction methods using lignocellulosic waste. *Biomass Conv Bioref* 2021. DOI: 10.1007/s13399-021-02062-z.

## Extraction, purification, and quantification of hesperidin from the immature *Citrus grandis/maxima* fruit Nepal cultivar

HEMANTH KUMAR MANIKYAM<sup>1,\*</sup>, PRATHIBHA TRIPATHI<sup>2</sup>, SANDEEP BALVANT PATIL<sup>3</sup>,  
JANARDAN LAMICHHANE<sup>4</sup>, MVNL CHAITANYA<sup>5</sup>, ABHINANDAN RAVSAHEB PATIL<sup>6</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science, North East Frontier Technical University. Aalo, West Siang 791001, Arunachal Pradesh, India.  
Tel./fax.: +91-3783-222-444, \*email: phytochem2@gmail.com

<sup>2</sup>Ayurveda Campus, Tribhuvan University. Kirtipur Rd, Kirtipur 44618, Nepal

<sup>3</sup>Dr. Shivajirao Kadam College of Pharmacy. Sangli Rd, Maharashtra 416301, India

<sup>4</sup>Department of Biotechnology, Organic farming and Natural Product Research Centre, Kathmandu University. Kavre, Dhulikhel 45200, Nepal

<sup>5</sup>Chitkara College of Pharmacy, Chitkara University. Punjab 140417, India

<sup>6</sup>Department of Pharmaceutics, Sanjay Ghodawat University. Kolhapur, Maharashtra 416118, India

Manuscript received: 9 April 2022. Revision accepted: 26 June 2022.

**Abstract.** Manikyam HK, Tripathi P, Patil SB, Lamichhane J, Chaitanya MVNL, Patil AR. 2022. Extraction, purification, and quantification of hesperidin from the immature *Citrus grandis/maxima* fruit Nepal cultivar. *Asian J Nat Prod Biochem* 20: 21-26. Hesperidin, a flavanone group of flavonoids and a 7-O-rutinoside of hesperitin, is abundant in citrus fruits. Pomelo or *Citrus grandis* (L.) Osbeck/ *C. maxima* (Burm.) Merr. is grown and cultivated in Nepal as seasonal edible fruit. Pomelo has flavanone and other chemical constituents. Flavanones like naringin, hesperidin, neohesperidin, and naringenin are present in the fruits of *C. grandis*. Many extraction techniques have been reported in the extraction and purification of hesperidin citrus fruits like sweet lemons, grapefruits, mandarins, etc. However, little information has been provided on hesperidin content in pomelo fruits. In order to estimate the quantity of hesperidin in the pomelo fruits of Nepal, we have first made a study. The immature pomelo fruits of size 2-7 cm were collected from the cultivation field and dried completely. The pulverized fruit powder with moisture content less than 15% was subjected to 5% methanolic acetic extraction and Dimethylformamide (DMF) solvent extraction in Soxhlet apparatus by refluxing at 90°C in 1:15 ratio in three subsequent extractions of each 1 hour. The concentrates were crystallized at freezing temperatures in methanol and further purified by repeating the crystallization process. The white amorphous powder was subjected to RP-HPLC analysis method, chromatographic conditions 285 nm wavelength, C18 column 4.6 mm x 15 cm x 5 µm, flow rate 1.2 mL/minute with injection volume 10 µL run time of 30 minutes, column temperature 40°C. Hesperidin yields were 0.12 g/100 g with acetic acid methanol with a purity of 87% and 0.17 g/100 g in DMF extraction with a purity of 90% with recoveries of 87% in acetic acid methanol extraction, and 90% in DMF extraction when compared with standard 93%. Further extraction does not yield any hesperidin content indicating a total of 0.15-0.17 g/100 g of hesperidin content in the whole immature fruit of *C. grandis/maxima*. Thus we found DMF and 5% methanolic acetic solvents as preferable solvents to extract hesperidin.

**Keywords:** Acetic acid, *Citrus grandis/maxima*, dimethylformamide, hesperidin, methanol, Nepal cultivar, RP-HPLC UV method

### INTRODUCTION

*Citrus grandis* (L.) Osbeck/ *C. maxima* (Burm.) Merr. commonly called pomelo (red pomelo), is an important member of the Rutaceae family, well known for its high morphological variability, nutritional and commercial values (Alam et al. 2014; Susandarini et al. 2020). It was found in Barbados and commercially grown in Morocco, Israel, Spain, South Africa, Brazil, the USA, and some Asian countries (Jokic et al., 2019). In Nepal, it is called Bhogate and is eaten as seasonal fruit. In Nepal, they make Bhogate sadheko, a common salad eaten by local people. In India commonly called Chakotra. China is the major producer of pomelo, which is more than 60% of the world's cultivation (Man et al., 2015). Many bioactive compounds like flavanones, flavones, flavonols, and anthocyanin are abundantly present in the pomelo fruit. Chemical ingredients like limonoid glycones and aglycones, coumarins, naringin, hesperidin flavanones, etc., are the most abundant compounds in pomelo fruits.

Naringin is the most abundant flavanone present, around 295 to 377 mg/L in grapefruits and pomelo. Many research studies have proved that the extracts of pomelo have a therapeutic effect on cancer, inflammation, diabetes, and neurodegenerative and cardiovascular diseases with powerful antioxidant activity (Damon et al. 1987; Ma et al. 2008). It was studied that active components in the pomelo extract by using high-performance liquid chromatography (HPLC) had naringin (11.90 ± 0.21 mg/g dried extract), hesperidin (12.04 ± 0.12 mg/g dried extract), neohesperidin (25.4 ± 0.12 mg/g dried extract), and naringenin (9.20 ± 0.19 mg/g dried extract) (El-Shafae and El-Domiatiy 2001; Lucker et al. 2002; Iglesias-Carres et al. 2019).

Hesperidin bio-flavanone, also known as hesperitin 7-rutinoside or 7-O-glycoside hesperitin (3,5,7 trihydroxyflavanone 7-rhamnoglucoside, C<sub>28</sub>H<sub>34</sub>O<sub>15</sub>). Figure 1, abundantly present in citrus fruits with a molecular weight of 610.57 with a melting point of 250-255°C and boiling point of 576.17°C, respectively. It is not soluble in water and highly soluble in alkaline solutions but

moderately soluble in propylene glycol and Poly ethylene glycol 400 (Kanaze et al. 2004). It is highly soluble in DMSO and DMF. Poorly soluble in solvents like ethanol, isopropanol, and methanol (when purified) (Pandey et al. 2015).

Hesperidin is generally considered safe for topical and systemic administrations, with 2% in topical applications; it has shown no adverse cutaneous reactions in mice for 9 days of administration (Xiong et al. 2019). In addition, the oral formulation Daflon-500 mg containing 10% hesperidin with a daily dosage of 100 mg showed no adverse effects in rats (Razavi and Hosseinzadeh 2019).

Multiple therapeutic effects of hesperidin include anti-inflammatory, lipid-lowering, analgesic, antioxidant, and anticancer activities. Other than mentioned health benefits, hesperidin is well-known to treat venous insufficiency, varicose veins, venous ulcers, and hemorrhoids (Paudyal and Haq, 2008; Man et al., 2019). By stimulating the release of the appetite-regulating hormone cholecystokinin (CCK) in enteroendocrine STC-1 cells, both hesperidin and hesperidin help treat obesity by controlling appetite (Magwaza et al., 2015). In one of the studies on the neuroblastoma cell line, hesperidin has exerted a neuroprotective effect against rotenone by its antioxidant effect and maintaining mitochondrial functioning (Wu et al. 2007).

In recent studies, attempted extraction of hesperidin from peels of *Citrus reticulata* Blanco mandarin using deep eutectic solvents (DESs), a sustainable green extraction technique using choline chloride: acetamide (1:2) and water DESs had shown most efficient extraction of hesperidin 112.14 mg/g (Hendrickson and Kesterson 1954). Extraction of hesperidin by ultrasound assistance has shown that solvent has a profound effect on the yields, along with an increase in temperature and extraction time (Kim and Lim 2020). The highest flavonoid yields were achieved by 58.4% ethanol concentration at 80°C, 40 mL/g solvents to feed concentration for 30 minutes, and hesperidin yields of 66.6%. The extraction conditions for the highest flavonoid yields based on a response surface methodology were 80.3°C, 58.4% (ethanol concentration), 40 mL/g (solvent/feed), and 30 min (Tamilselvam et al., 2013). Ethanol extraction was found to be more effective when compared to methanol (yield 57.3%) and acetone (yield 37.7%) (Tamilselvam et al., 2013). Some studies showed that the yield of hesperidin was higher in 70% methanol than in 80% ethanol at 35°C from mandarin peels (Caengprasath et al. 2013), and *C. sinensis* pulp was higher in 90% methanol than 90% ethanol at 55°C (Kim and Lim 2020).

In the present study for the first time, for the first time, we are extracting hesperidin from immature fruits of the *C. grandis/maxima* (pomelo) cultivar of Nepal and quantifying it.

## MATERIALS AND METHODS

**Plant material preparation:** *C. grandis/maxima* fruits were obtained from the local cultivation field Chisapani area, Kathmandu, Nepal, and immature fruits of size 2-7

cm see Figure 2, were dried till moisture content reached 15%. The 200 g of dried immature fruits of 2-7 cm were pulverized to 180 microns mesh size.

### Extraction of hesperidin using acetic acid methanol solvent

Weight separately 100 g of dried immature fruit powder of 2-7 cm size and place it in Soxhlet apparatus with 5% acetic in methanol (1:15 feed/solvent) and continued extraction for 1 hour at 100°C with a circulation of solvent. A total of 3 extractions were carried out at an interval of 1 hour with a 1:15 feed to solvent ratio consequently. The total extract is collected, filtered, and concentrated to 80% before chilling. The 80% reduced liquid concentrate was chilled at 5°C for a few hours and filtered. Finally, the precipitate of hesperidin is filtered and washed with pure methanol till a white-colored amorphous powder appears and is further quantified.

### Extraction of hesperidin with dimethylformamide

Weight separately 100 g of dried immature fruit powder of 2-7 cm size, place it in the maceration apparatus in DMF (1:15 feed/solvent), and continue stirring for 1 hour at 100°C with a solvent circulation. A total of 3 extractions were carried out at an interval of each 1 hour with a 1:15 feed to solvent ratio consequently. First, the extract is collected, filtered, and concentrated to 80% before chilling. Next, the 80% reduced liquid concentrate was chilled at 5°C for a few hours and filtered. Finally, the precipitate of hesperidin is filtered and washed with pure methanol till a white-colored amorphous powder appears and is further quantified.




Figure 1. Structure of hesperidin



Figure 2. Immature fruit (2-7 cm) of *Citrus grandis/maxima*

## Methods

### Loss on drying

Accurately weigh 1.0–2.0 g of sample, and spread it on a dry flat weighing bottle previously weighed. Dry the bottle at  $135 \pm 2^\circ\text{C}$  for 4 hours. Calculate loss on drying by the following formula, which should be less than 5.0%.

$$\text{Loss on drying\%} = (M1 - M2/M1 - M) \times 100\%$$

Where:

M : Tare weight of the weighing bottle (g)

M1 : Weight of the weighing bottle and sample before drying (g)

M2 : Weight of the weighing bottle and sample after drying (g)

### Assay (RP-HPLC)

#### Chromatographic system

**Chromatography system:** HPLC with UV detector; Column: 4.6 mm x 15 cm x 5  $\mu\text{m}$  packing C18; Wavelength: UV 284 nm; Flow rate: 1.2 mL/min; Column temperature:  $40^\circ\text{C}$ ; Injection volume: 10  $\mu\text{L}$ ; Run time: 30 minutes

**Mobile phase:** Methanol: water: Glacial acetic Acid: acetonitrile = 28:66:6:2 (V/V), diluent for sample preparation: dimethyl sulphoxide

#### Preparation of solution

**Test solution:** Accurately weigh 25 mg of sample to a 50 mL volumetric flask, dissolve and dilute to exactly 50 mL with dimethyl sulphoxide, and mix well.

**Reference solution:** Accurately weigh 25 mg of a hesperidin reference standard to a 50 mL volumetric flask, dissolve and dilute to exactly 50 mL with dimethyl sulphoxide, and mix well.

#### Test procedure

Inject reference solution, test solution, and record the chromatograms. Calculate the assay of hesperidin according to a standard external method based on an anhydrous basis. The result of the assay should be not less than 90%.

$$\text{Assay (\% w/w)} = (A_T/A_R \times W_S/W_R) \times P \times 100\%/1\text{-LOD}$$

Where:

$A_T$  : Main peak area in the chromatograms obtained with test solution;

$A_R$  : Main peak area in the chromatograms obtained with reference solution;

$W_S$  : Weight of sample, mg;

$W_R$  : Weight of reference standard, mg;

P : Assay of reference standard, %;

LOD : Loss on drying of sample, %.

### Method validation

Parameters like linearity, precision, accuracy, and limits of detection (LOD) and quantification (LOQ) were developed to determine the hesperidin content by the HPLC method. Calibration graphs for linearity were

determined using standard solution preparation of hesperidin at different concentrations of 20, 60, and 100  $\mu\text{g/mL}$ . The dilutions were injected in series, and the peak area was calculated for each concentration plotted against the peak area. By adding standard at three concentrations of 20, 60, and 100  $\mu\text{g/mL}$ , percentage recovery was calculated for each concentration to determine accuracy. Hesperidin was injected three times on the same day and on 3 different days to determine intra-day and inter-day variations to determine system precision. Both cases considered related standard deviation (RSD), a precision measure. By determining the LOD and LOQ, sensitivity was evaluated. The amount of analyte which gives a peak with a signal-to-noise ratio of 3 was defined as LOD, and the lowest amount of analyte with a signal-to-noise ratio of 10 was considered LOQ. The International Conference on Harmonization (ICH) was considered to validate the method.

### Quantification of hesperidin in concentrated methanolic acetic acid extract

The total methanolic acetic acid extracts thus collected after 3 subsequent extractions were combined and concentrated. The concentrated extract was then subjected to RP-HPLC chromatography as mentioned in the assay method for standard hesperidin, and the area of peak in correlation to the retention time of standard was analyzed. Finally, the amount of hesperidin was calculated from the calibration plot obtained by a regression equation.

## RESULTS AND DISCUSSION

Although hesperidin is a bioflavonoid in all citrus fruits, it is also known as Hesperitin 7-rutinoside or 7-O-glycoside hesperitin (3, 5, 7 trihydroxyfavanone 7-rhamnoglucoside). It is a highly economically viable phytochemical in demand for the Pharmaceutical, Nutraceutical, and cosmeceutical industries. It is a precursor molecule to synthesize Diosmin and a molecule used to treat neurological problems. Extraction of hesperidin using economically viable methods is needed in the present scenario looking at the abundant availability of raw material resources and market demand. Most immature citrus fruits varying in size from 2 mm- 3 mm contain 10-12% of hesperidin. Pomelo (*C. grandis/maxima*) is abundantly grown in Nepal as edible fruit. Pomelo fruit contains vitamin C, beta carotene, anthocyanin, hesperidin, and Naringin ingredients. Even though hesperidin content is found to be less in pomelo fruits, we attempted to quantify hesperidin in immature dried fruits for the first time in the Nepal cultivar of pomelo.

Many works have suggested using ethanol and methanol as solvents to extract hesperidin. In our present work, we have used two solvent methods, such as a 5% acetic acid in methanol and dimethylformamide, to extract hesperidin. The pH and temperature directly affect the recovery of the extraction of hesperidin. In experiments conducted by Hendrickson and Kesterson Florida horticulture society in 1955, they found hesperidin had

poor solubility in high-pH alcoholic solvents and high solubility in low-pH alcoholic solvents. Also, the optimal temperature for extraction should be from 90°C to 100°C (Alam et al., 2014). Research shows hesperidin solubility in DMF and DMSO (30 mg/mL) and soluble in ethanol and methanol (~1 mg/mL). Two separate experiments were carried out to extract hesperidin using 5% acetic acid in methanol and DMF in 1:15 feed to the solvent ratio for 3 hours consecutively at 100°C while stirring, and later the liquid extract was filtered and concentrated up to 80%. The concentrate was chilled at 5 °C for a few hours. Thus, the precipitate was filtered and purified by washing it with pure methanol. The obtained hesperidin was quantified using RP-HPLC UV analysis method, chromatographic conditions 285 nm wavelength, C18 column 4.6 mm x 15cm x 5 µm, flow rate 1.2 mL/minute with injection volume 10 µL run time of 30 minutes, column temperature 40°C. Hesperidin yields were 0.12 g/100g with acetic acid methanol with a purity of 87%, and 0.17 g/100 g in DMF extraction with a purity of 90% with recoveries of 87% in acetic acid methanol extraction, and 90% in DMF extraction when compared with standard 93% see Figure 3 and 4. Further extraction does not yield any hesperidin content indicating a total of 0.15-0.17 g/100 g of hesperidin content in the whole immature fruit of *C. grandis/maxima*. Thus we suggest using acetic acid methanol, methanol, or DMF as good solvents to extract high yields of hesperidin with good purity.

### Method validation

#### Linearity, LOD, and LOQ

It was confirmed that the linearity of the method for concentrations ranging from 20 to 100 µg/mL of hesperidin. Good linear regressions observed with calibration curves  $y = 0.3456x + 0.1573$   $R^2 = 0.9973$  (Figure 5 and Table 1). The minimum amount of analyte plant extract LOD used for detection and LOQ was 6.66 µg/mL and 20.19 µg/mL, with a retention time of 7.45 minutes.

#### Precision and accuracy

The precision and accuracy tests were performed by injecting the hesperidin sample three times within three consecutive days. Inter-day and Intra-day precision by RSD of the hesperidin peak areas ranged from 0.5-2.1, which is 2.5% under the limit as per ICH guidelines. By calculating the recovery %, accuracy was determined, and for hesperidin, it ranged between 89.99-102%, and the method was proven accurate (Table 2).


**Table 1.** Calibration data for the proposed HPLC method

Concentration (µg/mL)	Area (10 <sup>5</sup> mAU)
20	7.1
40	13.89
60	21.08
80	28.25
100	33.47
120	42.3


**Table 2.** Recovery and standard deviation data by HPLC

Compound	Amount added (µg/mL)	Amount recovered (µg/mL) <sup>a</sup>	Recovery (%) <sup>a</sup>	RSD (%)	
				Intra day	Inter day
Hesperidin	40	37.09 ± 0.26	92.75 ± 0.26	1.73	1.879
	80	81 ± 0.26	101.25 ± 0.26	2.12	0.156
	100	98.9 ± 1.69	98.9 ± 1.69	0.65	0.48


Note: <sup>a</sup> Mean ± SD (n=3) mean the sample was analyzed three times; <sup>b</sup> Samples were analyzed three times a day; <sup>c</sup> Sample was analyzed once a day over three consecutive days



**Figure 3.** Chromatogram of hesperidin standard 93%



**Figure 4.** Chromatogram of hesperidin (90% pure) extracted from the dried whole immature fruit of *Citrus grandis/maxima*



**Figure 5.** Linearity of hesperidin standard

## Discussion

Hesperidin flavanone glycoside is usually found in citrus fruits, and its aglycone is called hesperitin. Plants produce it as a defense mechanism. Different extraction techniques were used earlier to isolate hesperidin with significant yields from different varieties of citrus fruits. It has been found that classical solvent extraction of hesperidin is preferred. The highest flavonoid yield was achieved by 58.4% ethanol concentration at 80°C, 40 mL/g solvents to feed concentration for 30 minutes, and hesperidin yields of 66.6%. The extraction conditions for the highest flavonoid yields based on a response surface methodology were 80.3°C, 58.4% (ethanol concentration), 40 mL/g (solvent/feed), and 30 min (Tamilselvam et al., 2013). Ethanol extraction was found to be more effective when compared to methanol (yield 57.3%) and acetone (yield 37.7%) (Tamilselvam et al., 2013). Some studies showed that the yield of hesperidin was higher in 70% methanol than in 80% ethanol at 35°C from mandarin peels (Caengprasath et al. 2013), and *C. sinensis* pulp was higher in 90% methanol than 90% ethanol at 55°C (Kim and Lim 2020). In our present work, we first time tried to extract

hesperidin and quantify it using DMF and 5% methanolic acetic acid as solvents from the whole dried immature fruit of *C. grandis/maxima* of the Nepal cultivar variety. Two separate experiments were carried out to extract hesperidin using 5% acetic acid in methanol and DMF in 1:15 feed to the solvent ratio for 3 hours consecutively at 100°C while stirring, and later the liquid extract was filtered and concentrated up to 80%. The concentrate was chilled at 5°C for a few hours. Thus, the precipitate was filtered and then purified by washing it with pure methanol. The obtained hesperidin was quantified using RP-HPLC UV analysis method, chromatographic conditions 285 nm wavelength, C18 column 4.6 mm x 15 cm x 5 µm, flow rate 1.2 mL/minute with injection volume 10 µL run time of 30 minutes, column temperature 40°C. Hesperidin yields were 0.09 g/100 g with acetic acid methanol with a purity of 87% and 0.17 g/100 g in DMF extraction with a purity of 90% with recoveries of 87% in acetic acid methanol extraction, and 90% in DMF extraction, when compared with standard 93%, see Figures 3 and 4. It was confirmed that the linearity of the method for concentrations ranging from 20 to 100 µg/mL of hesperidin. Good linear regressions observed with calibration curves  $y = 0.3456x + 0.1573$ ,  $R^2 = 0.9973$  (Figure 5 and Table 1). The minimum amount of analyte plant extract LOD used for detection and LOQ was 6.66 µg/mL and 20.19 µg/mL, with a retention time of 7.45 minutes. The precision and accuracy tests were performed by injecting the hesperidin sample three times within the same and three consecutive days. Inter-day and Intra-day precision by RSD of the hesperidin peak areas ranged from 0.5-2.1, which is 2.5% under the limit as per ICH guidelines. By calculating the recovery %, accuracy was determined, and for hesperidin, it ranged between 89.99-102%, and the method was proven accurate (Table 2).

Further extraction does not yield any hesperidin content indicating a total of 0.15-0.17 g/100 g of hesperidin content in the whole immature fruit of *C. grandis/maxima*. Thus, we suggest using acetic acid methanol, methanol, or DMF

as good solvents to extract high yields of hesperidin with good purity.

## REFERENCES

- Alam MA, Subhan N, Rahman MM, Uddin SJ, Reza HM, Sarker SD. 2014. Effect of Citrus flavonoids, naringin and naringenin, on metabolic syndrome and their mechanisms of action. *Adv Nutr* 5 (4): 404-417. DOI: 10.3945/an.113.005603.
- Caengprasath N, Ngamukote S, Mäkynen K, Adisakwattana S. 2013. The protective effects of pomelo extract (*Citrus grandis* L. Osbeck) against fructose-mediated protein oxidation and glycation. *EXCLI J* 12: 491-502.
- Damon M, Flandre O, Michel F, Perdrix L, Labrid C, de Paulet AC. 1987. Effect of chronic treatment with a purified flavonoid fraction on inflammatory granuloma in the rat. Study of prostaglandin E2 and F2 $\alpha$  and thromboxane B2 release and histological changes. *Drug Res* 37 (10): 1149-1153.
- El-Shafae AM, El-Domiaty MM. 2001. Improved LC methods for the determination of diosmin and/or hesperidin in plant extracts and pharmaceutical formulations. *J Pharm Biomed Anal* 26 (4): 539-545. DOI: 10.1016/s0731-7085(01)00476-9.
- Hendrickson R, Kesterson JW. 1954. Recovery of Citrus Glucosides. *Proc Fla State Hort Soc* 67: 199-203.
- Iglesias-Carres L, Mas-Capdevila A, Bravo FI, Aragonès G, Muguera B, Arola-Arnal A. 2019. Optimization of a polyphenol extraction method for sweet orange pulp (*Citrus sinensis* L.) to identify phenolic compounds consumed from sweet oranges. *PLoS One* 14 (1): e0211267. DOI: 10.1371/journal.pone.0211267.
- Jokic S, Safranko S, Jakovljevic M, Cikoš A-M, Kajic N, Kolarevic F, Babic J, Molnar M. 2019. Sustainable green procedure for extraction of hesperidin from selected Croatian mandarin peels. *Processes* 7: 469. DOI: 10.3390/pr7070469.
- Kanaze FI, Kokkalou E, Georgarakis M, Niopas I. 2004. A validated solid-phase extraction HPLC method for the simultaneous determination of the citrus flavanone aglycones hesperetin and naringenin in urine. *J Pharm Biomed Anal* 36 (1): 175-181. DOI: 10.1016/j.jpba.2004.05.015.
- Kim DS, Lim SB. 2020. Extraction of flavanones from immature Citrus unshiu pomace: Process optimization and antioxidant evaluation. *Sci Rep* 10: 19950. DOI: 10.1038/s41598-020-76965-8.
- Lucker J, Tamer MK, Schwab W, Verstappen FWA, Plas LHWV, Bouwmesster HJ, Verhoeven HA. 2002. Monoterpene biosynthesis in lemon (*Citrus lemon*). *Eur J Biochem* 269: 3160-3171. DOI: 10.1046/j.1432-1033.2002.02985.x.
- Ma Y, Ye X, Hao Y, Xu G, Xu G, Liu D. 2008. Ultrasound-assisted extraction of hesperidin from Penggan (*Citrus reticulata*) peel. *Ultrason Sonochem* 15 (3): 227-232. DOI: 10.1016/j.ultsonch.2007.03.006.
- Magwaza LS, Opara UL, Cronje PJ, Landahl S, Ortiz JO, Terry LA. 2015. Rapid methods for extracting and quantifying phenolic compounds in citrus rinds. *Food Sci Nutr* 4 (1): 4-10. DOI: 10.1002/fsn3.210.
- Man G, Mauro TM, Y. Zhai Y, Kim PL, Cheung C, Hupe M, Crumrine D, Elias PM, Man M-Q. 2015. Topical hesperidin enhances epidermal function in an aged murine model. *J Invest Dermatol* 135: 1184-1187. DOI: 10.1038/jid.2014.486.
- Man M-Q, Yang B, Elias PM. 2019. Benefits of hesperidin for cutaneous functions. *Evid-Based Complement Alternat Med* 2019: 1-9. DOI: 10.1155/2019/2676307.
- Pandey RP, Gurung RB, Sohng JK. 2015. Dietary sources, bioavailability and biological activities of naringenin and its derivatives. In: Stacks NM (eds). *Apigenin and Naringenin: Natural Sources, Pharmacology and Role in Cancer Prevention*. Nova Science Publishers, Hauppauge, New York.
- Paudyal KP, Haq N. 2008. Variation of pomelo (*Citrus grandis* (L.) Osbeck) in Nepal and participatory selection of strains for further improvement. *Agrofor Syst* 72: 195-204. DOI: 10.1007/s10457-007-9088-z.
- Razavi BM, Hosseinzadeh H. 2019. A Review of the Effects of *Citrus grandis/maxima* (Grapefruit) and Its Flavonoids, Naringin, and Naringenin in Metabolic Syndrome, Bioactive Food as Dietary Interventions for Diabetes. (Second Edition). Academic Press. DOI: 10.1016/B978-0-12-813822-9.00034-5.
- Susandarini R, Subandiyah S, Daryono BS, Rugayah. 2020. Microsatellite polymorphism for molecular characterization of pomelo (*Citrus maxima*) accessions from Indonesia. *Biodiversitas* 21: 2390-2395. DOI: 10.13057/biodiv/d210608.
- Tamilselvam K, Braidy N, Manivasagam T, Essa MM, Prasad NR, Karthikeyan S, Thenmozhi AJ, Selvaraju S, Guillemin GJ. 2013. Neuroprotective effects of hesperidin, a plant flavanone, on rotenone-induced oxidative stress and apoptosis in a cellular model for Parkinson's disease. *Oxid Med Cell Longev* 2013: 102741. DOI: 10.1155/2013/102741.
- Wu T, Guan Y, Ya J. 2007. Determination of flavonoids and ascorbic acid in grapefruit peel and juice by capillary electrophoresis with electrochemical detection. *Food Chem* 100 (4): 1573-1579. DOI: 10.1016/j.foodchem.2005.12.042.
- Xiong H, Wang J, Ran Q, Lou G, Peng C, Gan Q, Hu J, Sun J, Yao R, Huang Q. 2019. Hesperidin: A therapeutic agent for obesity. *Drug Des Devel Ther* 13: 3855-3866. DOI: 10.2147/DDDT.S227499.